



Workshop on Multi-front Exotic phenomena in Particle and Astrophysics (MEPA 2022)

Time-varying resonant mass search

@

collider and beam dump experiments

Xiao-Ping Wang (王小平)

Beihang University (北航)

Dec, 17 @ MEPA2022

Base On: arXiv:2206.14221

Collaborated with: Jia Liu, Jinhui Guo, Yuxuan He and Kepan Xie






Dark Matter Evidence

- Evidence from different scale

Galaxy

Cluster


Universe





Dark matter mass range

Dark Sector Candidates, Anomalies, and Search Techniques



$$m_\phi \lesssim 30\text{eV} \left(\frac{250 \text{ km/s}}{\langle v^2 \rangle^{1/2}} \right)^{3/4} \left(\frac{\rho_{\text{DM}}}{0.4\text{GeV/cm}^3} \right)^{1/4}$$



Ultra light dark matter production

- Misalignment mechanism

$$\ddot{\phi} + 3H\dot{\phi} + m_{\phi}^2\phi = 0$$

- Classical Solution for wave-like dark matter

$$\phi(t) \approx \phi_1 \cos(m_{\phi}t) + \phi_2 \sin(m_{\phi}t)$$

- Dark matter local density

$$\rho_{\text{DM}} \approx \left(|\phi_1|^2 + |\phi_2|^2 \right) m_{\phi}^2$$

How to search for ultra light DM?



Dark matter-SM mediator




- Standard Model
= fermions + force

- Dark Sector
= DM + dark mediator

Standard Model of Elementary Particles

three generations of matter (fermions)			interactions / force carriers (bosons)	
I	II	III		
mass charge spin				
$\approx 2.2 \text{ MeV}/c^2$ $\frac{2}{3}$ $\frac{1}{2}$ u up	$\approx 1.28 \text{ GeV}/c^2$ $\frac{2}{3}$ $\frac{1}{2}$ c charm	$\approx 173.1 \text{ GeV}/c^2$ $\frac{2}{3}$ $\frac{1}{2}$ t top	0 0 1 g gluon	$\approx 125.09 \text{ GeV}/c^2$ 0 0 H higgs
$\approx 4.7 \text{ MeV}/c^2$ $-\frac{1}{3}$ $\frac{1}{2}$ d down	$\approx 96 \text{ MeV}/c^2$ $-\frac{1}{3}$ $\frac{1}{2}$ s strange	$\approx 4.18 \text{ GeV}/c^2$ $-\frac{1}{3}$ $\frac{1}{2}$ b bottom	0 0 1 γ photon	
$\approx 0.511 \text{ MeV}/c^2$ -1 $\frac{1}{2}$ e electron	$\approx 105.66 \text{ MeV}/c^2$ -1 $\frac{1}{2}$ μ muon	$\approx 1.7768 \text{ GeV}/c^2$ -1 $\frac{1}{2}$ τ tau	$\approx 91.19 \text{ GeV}/c^2$ 0 1 Z Z boson	
$< 2.2 \text{ eV}/c^2$ 0 $\frac{1}{2}$ ν_e electron neutrino	$< 1.7 \text{ MeV}/c^2$ 0 $\frac{1}{2}$ ν_μ muon neutrino	$< 15.5 \text{ MeV}/c^2$ 0 $\frac{1}{2}$ ν_τ tau neutrino	$\approx 80.39 \text{ GeV}/c^2$ ± 1 1 W W boson	
QUARKS			SCALAR BOSONS	
LEPTONS			GAUGE BOSONS VECTOR BOSONS	



This Talk

Phys.Rev.D 97 (2018) 4, 043001



Oscillating A' from wave-like DM

$$\mathcal{L} = -\frac{1}{4}F'_{\mu\nu}F'^{\mu\nu} + \frac{1}{2}m_0^2 A'_\mu A'^\mu + \epsilon e A'_\mu J_{\text{em}}^\mu$$

- Wave information to A'

$$\left(D_\mu \phi\right)^* D^\mu \phi \supset \left(g' Q_\phi\right)^2 \phi^* \phi A'_\mu A'^\mu$$

- Oscillating information of A'

$$\tilde{m}_0^2 = m_0^2 + \left(g' Q_\phi\right)^2 \left(\phi_1^* \phi_1 + \phi_2^* \phi_2 - \sqrt{\left(|\phi_1|^2 + |\phi_2|^2\right)^2 + \left(\phi_1 \phi_2^* + \phi_1^* \phi_2\right)^2} \right)$$

$$m_{A'}^2(t) = \tilde{m}_0^2 \left[1 + \left(2 \left(g' Q_\phi\right)^2 \frac{\sqrt{\left(|\phi_1|^2 + |\phi_2|^2\right)^2 + \left(\phi_1 \phi_2^* + \phi_1^* \phi_2\right)^2}}{\tilde{m}_0^2} \right) \cos^2 \left(m_\phi t\right) \right]$$

$\mathcal{K}$



Oscillating A' from wave-like DM

$$\mathcal{L} = -\frac{1}{4}F'_{\mu\nu}F'^{\mu\nu} + \frac{1}{2}m_0^2 A'_\mu A'^\mu + \epsilon e A'_\mu J_{\text{em}}^\mu$$

- Wave information to A'

$$\left(D_\mu \phi\right)^* D^\mu \phi \supset \left(g' Q_\phi\right)^2 \phi^* \phi A'_\mu A'^\mu$$

- For simply connect to UV model

$$\arg[\phi_1] = \arg[\phi_2] \text{ or } \phi_2 = 0$$



$$\tilde{m}_0 = m_0, \quad \kappa \equiv \frac{2 \left(g' Q_\phi\right)^2 \rho_{\text{DM}}}{m_\phi^2 m_0^2}$$

$$m_{A'}^2(t) = m_0^2(1 + \kappa \cos^2(m_\phi t))$$




Oscillating A' from wave-like DM

- Oscillating property

$$m_{A'}^2(t) = m_0^2(1 + \kappa \cos^2(m_\phi t))$$

$$= m_{A'}^2(t + \tau)$$



- Event number in the i th mass bin

$$N_i = \sigma_{\text{res}}^{(i)} \epsilon_i L \frac{\Delta t_i}{t_{\text{exp}}} = \frac{\sigma_{\text{res}}^{(i)} \epsilon_i L}{\tau} \int_{m_i}^{m_{i+1}} \left| \frac{dt}{dm_{\text{res}}} \right| dm_{\text{res}}$$


$$\frac{\tau}{m_0} f(y)$$



Oscillating A' from wave-like DM

- Oscillating property

$$f(y) = \frac{2y}{\pi \sqrt{(y^2 - y_{\min}^2)(y_{\max}^2 - y^2)}}$$




$$\frac{\int_{y_{\max}-\Delta}^{y_{\max}} f(y) dy}{\int_{y_{\min}}^{y_{\min}+\Delta} f(y) dy} \rightarrow \sqrt{\frac{y_{\max}}{y_{\min}}}$$



Oscillating A' effect on experiments?

- A kinetic mixing dark photon A' with $U(1)'$ interaction

$$\mathcal{L} = -\frac{1}{4}F'_{\mu\nu}F'^{\mu\nu} + \frac{1}{2}m_0^2 A'_\mu A'^\mu + \epsilon e A'_\mu J_{\text{em}}^\mu$$








Prompt dilepton final states

- General dilepton process

$$e^+e^- \rightarrow \gamma A', A' \rightarrow e^+e^-/\mu^+\mu^-$$



- BaBar data PRL 113, 201801 (2014)





Recast BaBar experiment result

PRL 113, 201801 (2014)



- Log likelihood ratio (LLR)

$$\text{LLR} = -2 \log \left[\frac{\text{Max}_{\vec{a}'} \prod_i \mathcal{N} \left(B_i - B(m_i, \vec{a}') - Sf_G(m_i) \mid B_i \right)}{\text{Max}_{\vec{a}} \prod_i \mathcal{N} \left(B_i - B(m_i, \vec{a}) \mid B_i \right)} \right]$$



Recast BaBar experiment result



- Traditional single-peak analysis

$$f_G(m_i) = \mathcal{N}(m_{A'} - m_i | \sigma_{\text{re}}^2)$$

- Double-peak analysis


$$f_S(m_i) = \int_{m_{\min}}^{m_{\max}} f\left(\frac{m'}{m_0}\right) \mathcal{N}(m_i - m' | \sigma_{\text{re}}^2) dm'$$

- Recast result:





Beam Dump Experiments




- General event number:

$$N(\epsilon, m_{A'}) = N_e \mathcal{C}' \epsilon^2 \frac{m_e^2}{m_{A'}^2} e^{-a_1 L_{sh} \Gamma_{A'}} (1 - e^{-a_2 L_{dec} \Gamma_{A'}})$$

- Including oscillation effect


$$N(\epsilon, m_0, \kappa) = \frac{1}{\tau} \int_{m_0}^{\sqrt{1+\kappa} m_0} N(\epsilon, m_{A'}) \left| \frac{dt}{dm_{A'}} \right| dm_{A'}$$

- Our recast result with $\kappa = 0$





Double-peak analysis result






Improvement with time-varying

- Oscillating property

$$m_{A'}^2(t) = \tilde{m}_0^2(1 + \kappa \cos^2(m_\phi t))$$




- Background event number

$$N_i^{\text{red}} = N_i \frac{1}{\tau} \int_{m_i}^{m_{i+1}} \left| \frac{dt}{dm_{A'}} \right| dm_{A'}$$



Improvement with time-varying

- Result with time-varying






Justify TDM with CMS open data

- CMS open data 2012

$$pp \rightarrow \ell^+ \ell^-, 8\text{TeV}, \sim 10 \text{ fb}^{-1}$$

- Luminosity is not constant





CMS open data

- Oscillating A' property

$$N_i = \frac{\sigma_{\text{res}}^{(i)} \epsilon_i L}{\tau} \int_{m_i}^{m_{i+1}} \left| \frac{dt}{dm_{\text{res}}} \right| dm_{\text{res}} \left[\frac{\tau}{\tilde{m}_0} f(y) \right]$$

- CMS differential event number

$$\frac{d\sigma_S}{dm_{\ell\ell}} = \epsilon_S \sigma_0 \times \delta(m_{\ell\ell} - m_{A'}(t))$$


$$\frac{dN_S}{dm_{\ell\ell}} = \int_{t_1}^{t_2} dt \mathcal{L}(t) \frac{d\sigma_S}{dm_{\ell\ell}} = \epsilon_S \sigma_0 \times \frac{2m_{\ell\ell}}{\pi \sqrt{m_{\ell\ell}^2 - m_0^2} \sqrt{(1+\kappa)m_0^2 - m_{\ell\ell}^2}} \frac{\tau_{A'}}{2} \left[\sum_{i^+} \mathcal{L}(t_i^+) + \sum_{i^-} \mathcal{L}(t_i^-) \right]$$

$$L \times \bar{f}(m_{\ell\ell}/m_0)$$



CMS open data

- Double-peak property



We can use DPM for CMS open data!





CMS result with time information

- Double-peak Method

$$\frac{dN_S}{dm_{\ell\ell}} = \sigma(m_{\ell\ell}) \epsilon_S \frac{f(m_{\ell\ell}/m_0)}{m_0} \times \frac{\tau_{A'}}{2} \sum_{i^\pm} \mathcal{L}(t_{i^\pm})$$

- Time-dependent Method

$$S_{ij} = \int_{t_i}^{t_i+\Delta t} dt \int_{m_j}^{m_j+\Delta m_{\ell\ell}} dm_{\ell\ell} \frac{1}{\sqrt{2\pi}\sigma_m} e^{-\frac{(m_{\ell\ell} - m_{A'}(t))^2}{2\sigma_m^2}} \times \mathcal{L}(t) \times \epsilon_S(m_{\ell\ell}) \sigma_0(m_{\ell\ell})$$





Summary

- **Time-varying resonant mass lead to double-peak feature in the invariant mass spectrum**
- **For mass around tens of MeV, the already excluded muon $(g - 2)\mu$ solution from A' becomes viable**
- **Time dependent method can improve the experiment sensitivity by a few orders**
- **We use the real collider data for a time-dependent resonance search and justify that our method works as expected**

Thank you!