The 32nd Texas Symposium on Relativistic Astrophysics



Contribution ID: 193

Type: Contributed talk in mini symposium

Alleviating H0-tension in T-dual Friedmann cosmologies

Friday, 15 December 2023 09:50 (10 minutes)

With increasing significance, the H0-tension between the Local Distance Ladder and Planck Λ CDM-analysis of the CMB points to a new principle of cosmological spacetime. Distinct from conventional extrapolations of general relativity in the Solar system, cosmological spacetime contains heat as a relic of the Big Bang based on Planck scale structures in black hole spacetimes. A path integral formulation produces a system of equations that satisfy T-duality in the Friedmann scale factor - a new symmetry in the Hubble expansion H(z) \simeq $(1+(6/5)\text{OmegaM0}[(1+z)^5-1])^{1/2}(1+z)$. It satisfies H0 \simeq 73 km/s/Mpc, q0 \simeq -1, OmegaM0 \simeq 0.25 relevant to tensions in (H0,q0,S8). [Based on van Putten, 2021, Phys. Lett. B, 823, 13637; 2020, MNRAS 491, L6.]

Primary author: VAN PUTTEN, maurice (INAF-OAS, Sejong University)

Presenter: VAN PUTTEN, maurice (INAF-OAS, Sejong University) **Session Classification:** Cosmology with large-scale structure