The 32nd Texas Symposium on Relativistic Astrophysics



Contribution ID: 300

Type: Poster

## Interacting Kilonovae: Long-lasting Electromagnetic Counterparts to Binary Mergers in the Accretion Disks of Active Galactic Nuclei

*Tuesday, 12 December 2023 17:27 (1 minute)* 

We investigate the dynamics and electromagnetic (EM) signatures of neutron star-neutron star (NS–NS) or neutron star-black hole (NS–BH) merger ejecta that occur in the accretion disk of an active galactic nucleus (AGN). Since most of the radiation energy has converted from the kinetic energy of merger ejecta, we call such an explosive phenomenon an interacting kilonova (IKN). It should be emphasized that IKNe are very promising, bright EM counterparts to NS–NS/BH–NS merger events in AGN disks. The peak luminosity of an IKN can be brighter than

 $10^{44}$  erg  $\cdot$  s<sup>-1</sup>, with similarity to superluminous supernovae (SLSNe) and tidal disruption events (TDEs). So the IKN could be one of the most energetic stellar optical transients in the universe. One can observe that the UV-optical-IR emission of an IKN can exceed the AGN background. This makes an IKN a very promising EM counterpart of GW events in the AGN disk for observation.

Primary author: REN, jia (nanjing university)

Co-authors: Prof. DAI, Zi-Gao (USTC); Mr CHEN, ken (nanjing university); Dr WANG, yun (PMO)

Presenter: REN, jia (nanjing university)

Session Classification: Accretion Processes