

# 第九届中国 LHC 物理年会 The 9th China LHC Physics Workshop (CLHCP2023)

Wednesday, 15 November 2023 - Monday, 20 November 2023

Shanghai

## Book of Abstracts



# Contents

$\Xi_c - \Xi'_c$ mixing From Lattice QCD . . . . .	1
Transverse-Momentum-Dependent Wave Functions of Pion from Lattice QCD . . . . .	1
Factorization of Non-Global LHC Observables and Resummation of Super-Leading Logarithms . . . . .	1
Constraint for a light charged Higgs boson and its neutral partners from top quark pairs at the LHC . . . . .	2
Automated calculation of Jet fragmentation at NLO in QCD . . . . .	2
Probing the spin-dependent fragmentation function in unpolarized pp and AA collisions at the LHC (Remote) . . . . .	3
Nonperturbative fitting in resummation calculation . . . . .	3
Top pair production in association with a W boson at the LHC . . . . .	3
Production and Decay of Top Quarks at Lepton Colliders at N3LO in QCD . . . . .	4
Operators Correlation in Electroweak Scattering at LHC . . . . .	4
Upgrade of Chinese LHCb Tier1 Site . . . . .	5
Probing positivity at LHC with exclusive photon-fusion processes . . . . .	5
Possible large CP violation in charmed Lambda_b decays . . . . .	5
Double-mixing CP violation . . . . .	6
Higgs boson pair production and decay to $b\bar{b}\gamma\gamma$ at NLO in QCD . . . . .	6
Higgs properties and new physics beyond the SM . . . . .	7
Partial NLO electroweak corrections to Higgs pair production in gluon fusion . . . . .	7
Probing Inelastic Dark Matter at the LHC, FASER and STCF . . . . .	7
Hybrid Renormalization for Quasi Distribution Amplitudes of A Light Baryon . . . . .	8
B meson anomalies and large $B^+ \rightarrow K^+\nu\bar{\nu}$ in non-universal $U(1)'$ models . . . . .	8
Resolving negative cross section of quarkonium hadroproduction using soft gluon factorization . . . . .	9

Cosmological Phase Transitions in Composite Higgs Models . . . . .	9
Theoretical Motivations for Hidden Light Bosons . . . . .	9
Neutrino as a window to TeV physics: from LHC to low-energy experiments . . . . .	10
Dark photon effects with the kinetic and mass mixing in $Z \rightarrow \tau^- \tau^+$ . . . . .	10
Probing the Higgs trilinear self-coupling through Higgs+jet production . . . . .	11
$B_{(s)} \rightarrow D_{(s)}^{**}$ form factors in HQEFT and model independent analysis of relevant semileptonic decays with NP effects . . . . .	11
Muon collider signatures for a $Z'$ with a maximal $\mu - \tau$ coupling in $U(1)_{L_\mu - L_\tau}$ . . . . .	11
Precise prediction for the top quark width . . . . .	12
Lattice study of singlet-assisted electroweak phase transition . . . . .	12
Observation of new structure in the $J/\psi/\psi$ mass spectrum in proton-proton collisions at $\sqrt{s} = 13$ TeV . . . . .	12
Electroweak sphalerons, scalar multiplets, and symmetry breaking patterns . . . . .	13
Search for Higgs boson pair production in the $b\bar{b}\mu\mu$ final state at the LHC (Remote) . . . . .	13
MatchingEFT.jl: An Automated Tool for Tree and One-Loop Level Matching . . . . .	14
Elliptic anisotropy of hard probes from parton scatterings in small collision systems . . . . .	14
Improved Asymptotic Formulae for Statistical Interpretation Based on Likelihood Ratio Tests . . . . .	15
Limiting FCNC induced by a CP symmetry of order 4 . . . . .	15
Aiming for Tops of ALPs with a Muon Collider . . . . .	15
The installation progress of upgrade Upstream Tracker at LHCb Upgrade II . . . . .	16
Explaining the CDF $W$ -mass shift and $(g - 2)_\mu$ in a $Z'$ scenario and its implications for the $b \rightarrow s \ell \ell^+ \ell^-$ processes . . . . .	16
Massive Scattering Amplitudes for Standard Model: On-shell Massless-Massive Correspondence . . . . .	16
The Prototype Design of PEB - a Component of the HGTD In-detector Electronics for the ATLAS Phase-II Upgrade . . . . .	17
Dark matter distribution, structure formation and the potential to distinguish thermal histories of dark matter. . . . .	17
Double Parton Scattering Effect in the Measurement of $W$ -Mass . . . . .	18
Hunting for sterile neutrino at colliders . . . . .	18
Probing Neutral Triple Gauge Couplings at the LHC, CEPC and SPPC . . . . .	18
$B \rightarrow D$ form factors beyond leading power and extraction of $ V_{cb} $ and $R(D)$ . . . . .	19

Unveiling time-varying signals of ultralight bosonic dark matter at collider and beam dump experiments . . . . .	19
Heavy long-lived coannihilation partner from inelastic Dark Matter model and its signatures at the LHC . . . . .	19
利用有限温度有效理论研究电弱相变 . . . . .	19
BBN 和 CMB 限制低能标新物理 . . . . .	19
Testing Complex Singlet Scalar Cosmology at the Large Hadron Collider . . . . .	19
Massive gauge theory with quasigluon for hot $SU(N)$ : Phase transition and thermodynamics . . . . .	20
Electroweak phase transitions proceed via bubbles vs. domain walls (Remote) . . . . .	20
Theory overview: Collider Physics . . . . .	20
Precision Higgs measurements at the LHC . . . . .	20
Search for HH production at the LHC . . . . .	20
BSM/rare Higgs . . . . .	20
Theory overview: Heavy flavor physics . . . . .	21
重味谱学 . . . . .	21
Search for rare decays at LHCb . . . . .	21
ALICE 实验中集体运动研究进展 . . . . .	21
Theory overview: QCD . . . . .	21
SM measurements at the LHC (+Top) . . . . .	21
CPV . . . . .	22
CEPC review . . . . .	22
ATLAS upgrade . . . . .	22
CMS upgrade . . . . .	22
LHCb upgrade . . . . .	22
ALICE upgrade . . . . .	22
Theory overview: Heavy ion physics . . . . .	23
ALICE 实验中奇异强子的产生 . . . . .	23
ALICE 实验中重夸克的产生与输运 . . . . .	23
Heavy Ion physics at ATLAS and CMS . . . . .	23
Theory Overview: Neutrino Physics . . . . .	23

SUSY results at LHC . . . . .	23
Dark matter and unconventional searches at the LHC . . . . .	23
Exotics (non-SUSY) searches at the LHC (+Flavor) . . . . .	24
Speech from SJTU-TDLI and CPS . . . . .	24
Speech from Funding Agency . . . . .	24
Speech from CPS High Energy Branch . . . . .	24
Announcements from LOC . . . . .	24
ATLAS report . . . . .	24
CMS report . . . . .	24
LHCb report . . . . .	24
ALICE report . . . . .	25
Theory overview: New Physics at the LHC: SMEFT and Beyond . . . . .	25
Machine Learning at LHC . . . . .	25
CLHCP2023 Summary . . . . .	25
Announcements . . . . .	25
Status of USTC-IME Pre-production sensor for the ATLAS High Granularity Timing Detector . . . . .	25
First results of USTC AC-coupled LGADs . . . . .	26
Electronics test results of Altiroc2 and Altiroc3 . . . . .	26
HGTD module assembly and module test at USTC . . . . .	26
ATLAS High granularity timing detector activity at IHEP/NJU . . . . .	26
IHEP-IME radiation hard LGAD sensor pre-production for ATLAS High granularity timing detector . . . . .	26
The Prototype Design of PEB - a Component of the HGTD In-detector Electronics for the ATLAS Phase-II Upgrade . . . . .	26
R&D of AC-LGAD based 4D tracker with precision timing information for HL-LHC and the future colliders . . . . .	26
Module performance study for the High Granularity Timing Detector of ATLAS . . . . .	27
Radiation Tolerance of the MUX64 for the High Granularity Timing Detector of ATLAS . . . . .	27
Automatic Module assembly development for ATLAS High granularity timing detector . . . . .	27
HGTD PEB DC/DC Power Block in Low Temperature and Magnetic Field Operation . . . . .	27

Performance of USTC-IME sensors in test beams at DESY and CERN . . . . .	27
CMS Endcap Timing Layer of MTD . . . . .	27
MTD sensor and assembly . . . . .	28
HGCal MAC status at IHEP . . . . .	28
HGCal SQC at IHEP and domestic silicon sensor R&D for calorimeter . . . . .	28
Computing vision QC in HGCal production . . . . .	28
基于 SystemC 对 FOCAL-E 像素层进行模拟分析 . . . . .	28
FOCAL 探测器像素层数据分析现状 . . . . .	28
ECAL simulation for LHCb Upgrade II . . . . .	28
Testbeam results of the LHCb Upgrade II ECAL prototype . . . . .	29
Performance of the GAGG crystal for LHCb Upgrade II ECAL . . . . .	29
ATLAS ITk Strip Module Site Qualification at IHEP . . . . .	29
Cold Noise Study in ATLAS ITk Strip Detectors . . . . .	29
SEE test of production ASICs in CSNS . . . . .	29
A study on the feasibility of CSNS becoming an ATLAS ITk sensor QA irradiation site . . . . .	29
Installation of LHCb Upstream Tracker . . . . .	29
Simulation of UT for LHCb Upgrade II . . . . .	30
Early Run3 $H \rightarrow \gamma\gamma$ fiducial Cross-Section measurement . . . . .	30
Test of CP Invariance in Higgs Boson Vector-Boson-Fusion Production Using the $H \rightarrow \text{gamgam}$ Channel with the ATLAS . . . . .	30
Novel photon energy calibration method and Higgs mass measurement . . . . .	30
(Remote) Measurement of the $H \rightarrow ZZ \rightarrow 4l$ cross-sections in pp collisions at 13.6 TeV with the ATLAS detector . . . . .	30
Measurements of inclusive and differential cross sections for the Higgs boson production and decay to four-leptons in proton-proton collisions at $\sqrt{s} = 13$ TeV . . . . .	30
Legacy search for the non-resonant production of Higgs boson pairs via gluon fusion and vector- bosonfusion in the $\text{XXXX}$ final state in proton-proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector (remote) . . . . .	30
Studies of new Higgs boson interactions through nonresonant HH production in the bbyy final state in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector . . . . .	31
Preliminary Results on Higgs Pair Production in Multi-lepton Channel with the ATLAS Experiment . . . . .	31
Search for diHiggs with VHH (Remote) . . . . .	31

Higgs mass and width measurement in ZZ to 4-leptons final state with full Run2 data (CMS)	31
Measurement of Higgs boson mass and width with LHC run2 data at the ATLAS experiment	31
B(s)->D(*)phi	31
Charm spectroscopy at LHCb	32
Interference fit to the double Jpsi mass spectrum at CMS	32
b-hadron FCNC decays at LHCb	32
Ds and D+ production in pPb	32
Psi(2S)/Jpsi versus multiplicity in pp	32
Bs->phiphi	32
lhcb EW results	32
Associated quarkonium production	33
Dicharm decay of beauty baryons	33
Omegac0 two-body hadronic decays at LHCb	33
B->D*K	33
Bs->Ds1K	33
Search for the exotic decay of the Higgs boson into a Z boson and a light pseudo-scalar decaying into two photons in pp collisions at 13TeV (CMS)	33
Search for extra Higgs bosons through same-sign top-quark production in association with an extra jet(Remote)	33
Searches for lepton flavour violation in Higgs boson decays, $H \rightarrow e\tau$ and $H \rightarrow \mu\tau$ , at ATLAS	34
Searches for Lepton-flavour-violating decays of the Higgs boson	34
Search for a standard model-like Higgs boson in the mass range between 70 and 110 GeV in the diphoton final state in proton-proton collisions at 13 TeV (CMS)	34
Evidence for the Higgs boson decay to a Z boson and a photon at the LHC	34
Run2 combination of the Higgs boson decay to a Z boson and a photon at the LHC	34
Interpretations of the measurements of Higgs boson production and decay rates and differential cross-sections based on the Nature paper(remote)	34
CMS Higgs and double Higgs combinations (Remote)	35
Measurement of ttW/ttH production in multilepton final states with ATLAS Run-2 data	35



Measurement of Higgs boson decaying into $b\bar{b}$ using Run2 dataset at the ATLAS experiment . . . . .	35
Non-prompt $\Lambda_c^+$ production with machine learning in p–Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV with ALICE . . . . .	35
ALICE 实验超氙产生测量 (remote) . . . . .	35
Study of heavy-flavor physics via semi-muonic decays with ALICE . . . . .	35
ALICE 实验轻核动量关联研究 (Remote) . . . . .	36
Measurement of $\Omega_c$ branching-fraction ratio at ALICE . . . . .	36
Pinning down the origin of collectivity in small systems with ALICE(Remote) . . . . .	36
Probing the nuclear structure with multiparticle correlation in Xe–Xe collisions . . . . .	36
基于 run3 数据的强子-奇异粒子关联研究现状 . . . . .	36
Measurement of inclusive $J/\Psi$ and $\Psi(2S)$ production at midrapidity in pp collisions at 13.6 TeV with ALICE . . . . .	36
Non-prompt $J/\Psi$ production in pp collisions with ALICE . . . . .	37
Measurement of inclusive $J/\Psi$ polarization at midrapidity in pp collisions at 13.6 TeV with ALICE . . . . .	37
$\Xi_{c0}$ production vs multiplicity via hadronic decay in pp at 13 TeV . . . . .	37
Study of strange hadron production in jets and the underlying events with ALICE . . . . .	37
ALICE 实验小系统碰撞中集体流研究 (Remote) . . . . .	37
ALICE 实验手征反常效应研究 (Remote) . . . . .	37
Measurement of the prompt and non-prompt $J/\Psi$ production in Pb–Pb collisions at 5.02 TeV with ALICE . . . . .	37
Study of jet quenching effects using jet-hadron correlations with ALICE(Remote) . . . . .	38
Study of beauty quark production properties via non-prompt D mesons with ALICE . . . . .	38
New perspectives on UV divergences of loops and the hierarchy problem . . . . .	38
[Cancel] Mono-X Signatures of an Absorbed Fermionic Dark Matter at the LHC . . . . .	38
Search for direct production of electroweakinos in final states with one lepton, jets and missing transverse momentum with the ATLAS detector . . . . .	38
Search for direct production of stau pairs in $\sqrt{s} = 13$ TeV $pp$ collisions with the ATLAS detector . . . . .	38
Search for lepton-flavour violation in high-mass dilepton final states using ATLAS run2 data . . . . .	39
Search for dark photons in rare Z boson decays with the ATLAS detector . . . . .	39

Search for a new $Z'$ gauge boson in the $\mu\mu\nu$ final state with the ATLAS detector . . . . .	39
A search for heavy right-handed Majorana neutrinos with ATLAS . . . . .	39
Progress of the RPC development towards the production for ATLAS Phase-II upgrade . . . . .	39
CMS-GEM upgrade progress and GEM assembly at Peking University . . . . .	39
CMS-GEM ME0 electronics board design, prototyping and production plan . . . . .	40
CMS iRPC Backend and Trigger status . . . . .	40
Data analysis for iRPC BE/TRG beam test . . . . .	40
Grid computing for LHC experiments in China . . . . .	40
Dark SHINE Simulation software framework . . . . .	40
IHEP ROC Status Report . . . . .	40
Run3 CMS EGamma performance (Remote) . . . . .	41
Optimal transport solutions for pileup mitigation at hadron colliders (Remote) . . . . .	41
Ultimate calibration and performance of the CMS Electromagnetic Calorimeter in LHC Run 2(Remote) . . . . .	41
RPC operation in the ATLAS experiment(Remote) . . . . .	41
Tau trigger scale factor studies . . . . .	41
CMS DQM-DC operations and performance during Run3 data-taking . . . . .	41
Constituent-based W-boson tagging with the ATLAS detector . . . . .	41
Performance of heavy flavour jet identification in boosted topologies with full Run2 data . . . . .	42
Observation of $WW\gamma$ production . . . . .	42
VBS $W\gamma$ at CMS . . . . .	42
$ZZ$ cross section measurement at 13.6TeV with the ATLAS detector . . . . .	42
Evidence of pair-production of longitudinally polarized vector bosons in $ZZ4l$ final states . . . . .	42
Observation of $WZ\gamma$ production in pp collisions with ATLAS detector . . . . .	42
Precise measurement of $Z\gamma$ +jets final state and search for neutral triple gauge couplings . . . . .	42
Observation and measurement of the cross-sections of the electroweak and total production of a $Z\gamma$ pair in association with two jets in pp collisions at $\sqrt{s} = 13\text{TeV}$ with the ATLAS detector (remote) . . . . .	43
Observation of Four-Top-Quark Production in the Multi-lepton Final State with the ATLAS Experiment (remote) . . . . .	43
Search for heavy Higgs bosons decaying to atop quark pair with the ATLAS detector . . . . .	43

Search for a new high-mass resonance decaying into $Z\gamma$ final state in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector . . . . .	43
Search for leptoquarks decaying into the $b\tau$ final state in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector . . . . .	43
Search for the production of an excited bottom quark decaying to $tW$ (leptonic top + hadronic $W$ ) in pp collisions at 13 TeV (CMS) . . . . .	44
Search for a new heavy boson $W'$ decaying to a top quark and a bottom quark with the ATLAS detector . . . . .	44
Search for $\bar{\nu}_\tau \nu_\tau \rightarrow \bar{\nu}_\tau \nu_\tau$ production in a final state 2 with one lepton or two opposite-sign leptons using the full Run 2 $\bar{\nu}_\tau \nu_\tau$ collisions data at $\sqrt{s} = 13$ TeV . . . . .	44
Observation and differential cross-section measurements of electroweak $\bar{\nu}_\tau \nu_\tau$ production in $\bar{\nu}_\tau \nu_\tau$ collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector (Remote) . . . . .	44
Measurements of energy correlators and alphas extraction . . . . .	44
First Direct Observation of Collider Neutrinos with FASER . . . . .	44
Study of isolated photon production with ALICE . . . . .	45
First Results from the Search for Dark Photons with the FASER Detector at the LHC . . . . .	45
Combination and summary of ATLAS dark matter searches interpreted in the context of 2HDM+a . . . . .	45
b-tagging trigger calibration with $t\bar{t}$ event . . . . .	45
Improved Asymptotic Formulae for Statistical Interpretation Based on Likelihood Ratio Tests . . . . .	45
Status of Progress of CEPC Calorimeter R&D . . . . .	45
CEPC Flavor Physics . . . . .	45
Silicon detector at the CEPC . . . . .	46
CEPC New Physics Studies . . . . .	46
Electroweak Physics at the CEPC . . . . .	46
Consideration on CEPC Electronics . . . . .	46
“小爆炸”：相对论重离子碰撞 . . . . .	46
Performance of ATLAS Photon ID . . . . .	46
Search for Higgs pairs in the $b\bar{b}\gamma\gamma$ final state with the ATLAS Experiment . . . . .	46
Performance and Calibration of quark/gluon jet tagger and search for new resonances with quark/gluon tagged dijet events . . . . .	47
Test results of Quality Control Test Structure of the USTC-IME LGAD pre-production . . . . .	47
Improvement of jet reconstruction in forward region with time information from HGTD. . . . .	47

ACP technique for flip chip bonding . . . . .	47
LGAD based Beam monitor development . . . . .	47
Dark Photon search initiative at SHINE facility (Dark SHINE) . . . . .	47
Dark SHINE Simulation software framework . . . . .	47
R&D of Dark SHINE LYSO Crystal ECAL . . . . .	48
R&D of Dark SHINE Strip Tracker . . . . .	48
R&D of Dark SHINE Hadronic Calorimeter (HCAL) . . . . .	48
Development of readout electronics for Dark SHINE ECAL . . . . .	48
Simulation study of the muon flux produced by 1.6 GeV electrons in a beam dump . . . .	48
Particle physics opportunities at the Shanghai SHINE facility with secondary beams . . .	48
Development of a muon spin polarization monitor for the J-PARC Muon g-2/EDM experiment . . . . .	49
Muon Lifetime Measurement with Run-1 data of Muon g-2 Experiment at Fermilab . . .	49
Development of a muon entrance trigger system for the PSI muEDM experiment . . . .	49
Performance of a prototype PSI muEDM muon entrance detector measured with test beam data . . . . .	49
Reconstruction of Neutrino Events in TRIDENT Project based on GNN . . . . .	49

**Theory / 1**

## $\Xi_c - \Xi'_c$ mixing From Lattice QCD

**Author:** Hang Liu<sup>one</sup>

**Corresponding Author:** liuhang303@sjtu.edu.cn

In heavy quark limit, the lowest-lying charmed baryons with two light quarks can form an SU(3) triplet and sextet. The  $\Xi_c$  in the SU(3) triplet and  $\Xi'_c$  in the sextet have the same  $J^{PC}$  quantum number and can mix due to the finite charm quark mass and the fact the strange quark is heavier than the up/down quark. We explore the  $\Xi_c$ - $\Xi'_c$  mixing by calculating the two-point correlation functions of the  $\Xi_c$  and  $\Xi'_c$  baryons from lattice QCD. Based on the lattice data, we adopt two independent methods to determine the mixing angle between  $\Xi_c$  and  $\Xi'_c$ . After making the chiral and continuum extrapolation, it is found that the mixing angle  $\theta$  is  $1.2^\circ \pm 0.1^\circ$ , which seems insufficient to account for the large SU(3) symmetry breaking effects found in weak decays of charmed baryons.

**Theory / 2**

## Transverse-Momentum-Dependent Wave Functions of Pion from Lattice QCD

**Author:** Minhuan Chu<sup>1</sup>

**Co-author:** Wei Wang

<sup>1</sup> *Shanghai Jiao Tong University*

**Corresponding Author:** tony\_chu@sjtu.edu.cn

We present a first lattice QCD calculation of the transverse-momentum-dependent wave functions (TMDWFs) of the pion using large-momentum effective theory. Numerical simulations are based on one ensemble with 2+1+1 flavors of highly improved staggered quarks action with lattice spacing  $a = 0.121$  fm from the MILC Collaboration, and one with 2 +1 flavor clover fermions and tree-level Symanzik gauge action generated by the CLS Collaboration with  $a = 0.098$  fm. As a key ingredient, the soft function is first obtained by incorporating the one-loop perturbative contributions and a proper normalization. Based on this and the equal-time quasi-TMDWFs simulated on the lattice, we extract the light-cone TMDWFs. The results are comparable between the two lattice ensembles and a comparison with phenomenological parametrization is made. Our studies provide a first attempt of ab initio calculation of TMDWFs which will eventually lead to crucial theory inputs for making predictions for exclusive processes under QCD factorization.

**Theory / 3**

## Factorization of Non-Global LHC Observables and Resummation of Super-Leading Logarithms

**Author:** Dingyu Shao<sup>1</sup>

<sup>1</sup> *Fudan University*

**Corresponding Author:** dyshao@fudan.edu.cn

We present a systematic formalism based on a factorization theorem in soft-collinear effective theory to describe non-global observables at hadron colliders, such as gap-between-jets cross sections. The cross sections are factorized into convolutions of hard functions, capturing the dependence on the partonic center-of-mass energy, and low-energy matrix elements, which are sensitive to the low

scale characteristic of the veto imposed on energetic emissions into the gap between the jets. The scale evolution of both objects is governed by a renormalization-group equation, which we derive at one-loop order. By solving the evolution equation for the hard functions for arbitrary  $2 \rightarrow M$  jet processes in the leading logarithmic approximation, we accomplish for the first time the all-order resummation of the so-called “super-leading logarithms” discovered in 2006, thereby solving an old problem of quantum field theory. We study the numerical size of the corresponding effects for different partonic scattering processes and explain why they are sizable for  $pp \rightarrow 2\text{jets}$  processes, but suppressed in  $H/Z + \text{jet}$  production. The super-leading logarithms are given by an alternating series, whose individual terms can be much larger than the resummed result, even in very high orders of the loop expansion. Resummation is therefore essential to control these effects. We find that the asymptotic fall-off of the resummed series is much weaker than for standard Sudakov form factors.

TeV / 5

## Constraint for a light charged Higgs boson and its neutral partners from top quark pairs at the LHC

**Authors:** Jun Gao<sup>one</sup>; 淳浩傅<sup>one</sup>

**Corresponding Author:** fuchh21@sjtu.edu.cn

The charged Higgs boson plays an essential role in distinguishing between a wide variety of standard model extensions with multiple Higgs doublets. We study the prospect of a light charged Higgs boson, produced by top quark pairs at the Large Hadron Collider (LHC), and decaying into a  $W$  boson and a pair of bottom quarks via an intermediate neutral Higgs boson ( $H_i$ ). We reinterpret the cross sections of  $WWbb\bar{b}\bar{b}$  final states measured by the ATLAS collaboration at LHC 13 TeV in the presence of the aforementioned decay, in a relatively wide range of Higgs masses. We find improved agreements with the data and obtain limits on the total branching ratio of the decay chain. The limits impose the strongest constraints on the parameter space of type-I two-Higgs-doublet model for most Higgs masses sampled when  $H_i$  is the  $CP$ -odd Higgs boson  $A$ . We also calculate potential constraints with pseudodata in high-luminosity runs of the LHC.

Theory / 6

## Automated calculation of Jet fragmentation at NLO in QCD

**Authors:** 重阳刘<sup>1</sup>; 晓民沈<sup>1</sup>; 斌周<sup>1</sup>; 俊高<sup>1</sup>

<sup>1</sup> 上海交通大学粒子物理与核物理研究所

**Corresponding Author:** liucy1999@sjtu.edu.cn

We present FMNLO, a framework to combine general-purpose Monte Carlo generators and fragmentation functions (FFs). It is based on a hybrid scheme of phase-space slicing method and local subtraction method, and accurate to next-to-leading order (NLO) in QCD. The new framework has been interfaced to MG5\_aMC@NLO and made publicly available in this work. We demonstrate its unique ability by giving theoretical predictions of various fragmentation measurements at the LHC, followed by comparison with the data. With the help of interpolation techniques, FMNLO allows for fast calculation of fragmentation processes for a large number of different FFs, which makes it a promising tool for future fits of FFs. As an example, we perform a NLO fit of parton fragmentation functions to unidentified charged hadrons using measurements at the LHC. We find the ATLAS data from inclusive dijet production show a strong constraining power. Notable disparities are found between our gluon FF and that of BKK, DSS and NNFF, indicating the necessities of additional constraints and data for gluon fragmentation function.

Theory / 7

## Probing the spin-dependent fragmentation function in unpolarized $pp$ and AA collisions at the LHC (Remote)

**Author:** Shu-Yi WEI<sup>1</sup>

<sup>1</sup> *Shandong University*

**Corresponding Author:** shuyi@sdu.edu.cn

The longitudinal spin transfer represents the probability density of producing longitudinally polarized hadrons from longitudinally polarized quarks or circularly polarized gluons. It thus was usually measured in polarized reactions or high-energy collisions where weak interaction dominates. In this work, we propose the dihadron polarization correlation as a novel probe of this quantity. Such an observable does not require the fragmenting partons to be polarized and therefore can be measured in the currently available experimental facilities, such as Belle, RHIC, Tevatron, and the LHC. We make quantitative predictions for these experiments. In light of the data already harvested, the experimental investigation of this observable provides more opportunity for the quantitative study of the longitudinal spin transfer. In particular, the measurements in  $pp$  collisions can significantly constrain the fragmentation function of a circularly polarized gluon.

### References

- [1] Probing the longitudinal spin transfer via dihadron polarization correlations in unpolarized  $e^+e^-$  and  $pp$  collisions, Hao-Cheng Zhang, Shu-Yi Wei, Phys. Lett. B 839 (2023) 137821.  
 [2] X. Li, Z.X. Chen, S. Cao, S.Y. Wei, arXiv:2309.09487 (2023).

Theory / 8

## Nonperturbative fitting in resummation calculation

**Authors:** Josh Isaacson<sup>1</sup>; Yao Fu<sup>2</sup>; C.-P. Yuan<sup>3</sup>

<sup>1</sup> *Fermi National Accelerator Laboratory*

<sup>2</sup> *University of Science and Technology of China*

<sup>3</sup> *Michigan State University*

**Corresponding Author:** yfu@mail.ustc.edu.cn

The resummation calculation (ResBos) is a widely used tool for the simulation of single vector boson production at colliders. As the improvement over the ResBos code by increasing the accuracy from NNLL+NLO to N<sup>3</sup>LL+NNLO, the nonperturbative function needs to be updated. We propose a new non-perturbative function (IFY) that includes information about the rapidity of the system. The IFY functional form was fitted to data from fixed target experiments, the Tevatron, and the LHC. We find that the non-perturbative function has mild rapidity dependence based on the results of the fit.

Theory / 9

## Top pair production in association with a W boson at the LHC

**Authors:** Bayu Hartanto Heribertus<sup>None</sup>; Fernando Febres Cordero<sup>None</sup>; Giuseppe Bevilacqua<sup>None</sup>; Jasmina Nasufi<sup>None</sup>; Laura Reina<sup>None</sup>; Malgorzata Worek<sup>None</sup>; Manfred Kraus<sup>None</sup>; 环宇毕<sup>1</sup>

<sup>1</sup> 北京大学

**Corresponding Author:** bihy@pku.edu.cn

This presentation provides a phenomenological analysis of the production of  $t\bar{t}W$  and  $t\bar{t}Wj$  in the context of the Standard Model at the LHC with a center-of-mass energy of  $\sqrt{s} = 13$  TeV. The analysis is focused on investigating various aspects of the computation, such as the impact of employing different theoretical modeling approaches, cross section ratio  $\sigma_{t\bar{t}W^+}/\sigma_{t\bar{t}W^-}$ , and the charge asymmetries.

**Theory / 10**

## Production and Decay of Top Quarks at Lepton Colliders at N3LO in QCD

**Authors:** Long Chen<sup>1</sup>; Xiang Cheng<sup>2</sup>; Xin Guan<sup>2</sup>; Yan-Qing Ma<sup>2</sup>

<sup>1</sup> Shandong University

<sup>2</sup> Peking University

**Corresponding Author:** longchen@sdu.edu.cn

In this talk we will discuss the recent processes in the calculation of the high-order perturbative corrections to the semi-inclusive production and decay of top quarks at lepton colliders at N3LO in QCD. In particular, the talk will be focusing on the first high-precision calculation of the complete QCD corrections to the top-quark decay width  $\Gamma_t$ ,  $W$ -helicity fractions and semi-inclusive distributions to the third order in the strong coupling constant  $\alpha_s$ . We find, in particular, that the pure  $\mathcal{O}(\alpha_s^3)$  correction decreases  $\Gamma_t$  by 0.8% of the previous  $\mathcal{O}(\alpha_s^2)$  result, exceeding considerably the error estimated by the usual scale-variation prescription. With this critical piece of correction included, we arrive at the to-date most precise theoretical prediction which meets the envisaged precision request by future hadron and lepton colliders.

**Theory / 11**

## Operators Correlation in Electroweak Scattering at LHC

**Author:** Jian-Nan Ding<sup>1</sup>

**Co-authors:** Fu-Sheng Yu <sup>2</sup>; Qing-Hong Cao <sup>1</sup>

<sup>1</sup> Peking University

<sup>2</sup> Lanzhou University

**Corresponding Author:** dingjn23@pku.edu.cn

To probe new physics without prior assumptions on UV models, the correlation of operators could be crucial in exposing the structure of UV completion.

When operators arise from the same heavy resonance, they are likely to correlate and their Wilson coefficients exhibit non-trivial relation, since both of them depend on the same UV parameters.

The aim of EFT analysis is to discover the correlation among operators, which might shed lights on UV completion.

For specific, if a precise measurement is consistent with the SM theory prediction, it might originate from a coherent cancellation among higher dimensional operators.

In this work, we investigate a strongly correlated cancellation of operators in electroweak scattering and attempt to expose the corresponding UV structure.

We also examine the operators correlation through a coupled channel analysis method and demonstrate that this correlation persists even when considering the uncertainties at HL-LHC.



On the other hand, since the operators connect different scattering channels through the reduction of  $H$  into  $v$  or  $h$ , the operators correlation in single top production will predict the total cross section of  $thq$  production, and the correlation in  $pp \rightarrow h\gamma$  can precisely examine the new physics effects of the indirect search on the weak magnetic moment  $a_W$ .

Theory / 12

## Upgrade of Chinese LHCb Tier1 Site

**Author:** Xiaowei JIANG<sup>1</sup>

<sup>1</sup> *IHEP*

**Corresponding Author:** jiangxw@ihep.ac.cn

The report will introduce the construction status of the first LHCb (WLCG) tier1 site in China. The content will refer the progress of requesting the tier1 site, the site scalability, the current status and future work. The site is built under the cooperation of Chinese LHCb Collobaration and IHEP computing center.

Theory / 13

## Probing positivity at LHC with exclusive photon-fusion processes

**Authors:** Chi Shu<sup>1</sup>; Jiayin Gu<sup>1</sup>

<sup>1</sup> *Fudan University*

**Corresponding Author:** cshu20@fudan.edu.cn

By tagging one or two intact protons in the forward direction, it is possible to select and measure exclusive photon-fusion processes at the LHC. The same processes can also be measured in heavy ion collisions, and are often denoted as ultraperipheral collisions (UPC) processes. Such measurements opens up the possibility to probe certain dimension-8 operators and their positivity bounds at the LHC. As an demonstration, we perform a phenomenological study on the  $\gamma\gamma \rightarrow \ell^+\ell^-$  processes, and find out that the measurements of this process at the HL-LHC provides reaches on certain dimension-8 operator coefficients that are comparable to the ones at future lepton colliders. We also point out that the  $\gamma q \rightarrow \gamma q$  process could potentially have better reaches on similar types of operators due to its larger cross section, but a more detailed experimental study is need to estimate the background of this process. The validity of effective field theory (EFT) and the robustness of the positivity interpretation are also discussed.

Theory / 14

## Possible large CP violation in charmed Lambda\_b decays

**Author:** Yin-Fa Shen<sup>1</sup>

**Co-authors:** Jian-Peng Wang<sup>2</sup>; Qin Qin<sup>3</sup>

<sup>1</sup> *HUST*

<sup>2</sup> *Lanzhou Univeristy*

<sup>3</sup> *Huazhong University of Science and Technology*

**Corresponding Author:** syf70280@hust.edu.cn

We propose that the cascade decay  $\Lambda_b \rightarrow D(\rightarrow K^+\pi^-)N(\rightarrow p\pi^-)$  may serve as the discovery channel for baryonic CP violation. This decay chain is contributed by dominantly the amplitudes with the intermediate  $D$  state as  $D^0$  or  $\bar{D}^0$ . The large weak phase between the two kinds of amplitudes suggests the possibility of significant CP violation. While the presence of undetermined strong phases may complicate the dependence of CP asymmetry, our phenomenological analysis demonstrates that CP violation remains prominent across a broad range of strong phases. The mechanism also applies to similar decay modes such as  $\Lambda_b \rightarrow D(\rightarrow K^+K^-)\Lambda$ . Considering the anticipated luminosity of LHCb, we conclude that these decay channels offer a promising opportunity to uncover CP violation in the baryon sector.

**Theory / 15**

## Double-mixing CP violation

**Author:** Yin-Fa Shen<sup>one</sup>

**Co-authors:** Qin Qin <sup>1</sup>; wen jie Song <sup>2</sup>

<sup>1</sup> *Huazhong University of Science and Technology*

<sup>2</sup> 华中科技大学

**Corresponding Author:** jaysong@hust.edu.cn

We propose a new kind of CP violation effect —the double-mixing CP asymmetry —in a type of cascade decays that involves at least two mixing neutral mesons in the decay chain. It is induced by the interference between different oscillation paths of the neutral mesons in the decay process. The double-mixing CP asymmetry is of critical importance for phenomenology, providing opportunities for clean determination of CKM phase angles free of uncertainties induced by the strong dynamics. To illustrate this point, we perform a phenomenological analysis on two examples:  $B_s^0 \rightarrow \rho^0 K \rightarrow \rho^0(\pi^-\ell^+\nu_\ell)$  and  $B^0 \rightarrow D^0 K \rightarrow D^0(\pi^+\ell^-\bar{\nu}_\ell)$ . Our results demonstrate that the double-mixing CP asymmetry can be numerically significant in the absence of strong phases, as shown by the former example. Additionally, the latter example showcases the direct extraction of weak and strong phases from data, without the need for theoretical inputs.

**TeV / 16**

## Higgs boson pair production and decay to $b\bar{b}\gamma\gamma$ at NLO in QCD

**Authors:** Dan Zhao<sup>1</sup>; Haitao Li<sup>2</sup>; Jian Wang<sup>3</sup>; Zongguo Si<sup>4</sup>; xiao zhang<sup>4</sup>

<sup>1</sup> 山东大学

<sup>2</sup> *haitao.li@sdu.edu.cn*

<sup>3</sup> *Shandong University*

<sup>4</sup> *shandong university*

**Corresponding Author:** zhaodan@mail.sdu.edu.cn

We calculate the total cross-section and differential distributions of Higgs boson pair production and decay to  $b\bar{b}\gamma\gamma$  at NLO in QCD.

Plenary Session / 17

## Higgs properties and new physics beyond the SM

**Author:** Bin Yan<sup>1</sup>

<sup>1</sup> *IHEP*

**Corresponding Author:** yanbin@ihep.ac.cn

The discovery of the Higgs boson at the Large Hadron Collider (LHC) has opened a new era in particle physics. Precise measurements of the properties of the Higgs boson are crucial for addressing several fundamental questions in the field. These include understanding the mechanism behind electroweak symmetry breaking, unraveling the origin of particle masses, and exploring potential sources of CP violation

that could explain the matter-antimatter asymmetry in the universe, and so on. In this talk, I will provide an overview of the recent advancements in Higgs physics, both within the framework of the Standard Model (SM) and beyond. By examining the latest research, we will gain insights into the properties and behavior of the Higgs boson, shedding light on the fundamental workings of the universe.

TeV / 18

## Partial NLO electroweak corrections to Higgs pair production in gluon fusion

**Authors:** Haitao Li<sup>1</sup>; Jian Wang<sup>2</sup>; Zongguo Si<sup>3</sup>; xiao zhang<sup>4</sup>; 丹赵<sup>5</sup>

<sup>1</sup> *haitao.li@sdu.edu.cn*

<sup>2</sup> *Shandong University*

<sup>3</sup> *shandong university*

<sup>4</sup> *ShanDong University*

<sup>5</sup> *山东大学*

**Corresponding Author:** zhaangx@mail.sdu.edu.cn

We calculated partial SM NLO electroweak corrections to Higgs pair production in gluon fusion, which are proportional to triple Higgs self-interacting constant. Then we did the same calculations with the Standard Model Effective Field Theory (SMEFT) dimension-6 Higgs self-interacting operators, and calculated the cross sections corresponding to different SMEFT parameters.

Theory / 19

## Probing Inelastic Dark Matter at the LHC, FASER and STCF

**Author:** Chih-Ting Lu<sup>1</sup>

**Co-authors:** Jianfeng Tu<sup>1</sup>; Lei Wu<sup>2</sup>

<sup>1</sup> *Nanjing Normal University*

<sup>2</sup> *Nanjing Normal Univeristy*

**Corresponding Author:** ctlu@nju.edu.cn

In this talk, we explore the potential of probing the inelastic dark matter (DM) model with an extra  $U(1)_D$  gauge symmetry at the Large Hadron Collider, ForwArd Search ExpeRiment and Super Tau Charm Factory. To saturate the observed DM relic density, the mass splitting between two light dark states has to be small enough, and thus leads to some distinctive signatures at these colliders. By searching for the long-lived particle, the displaced muon-jets, the soft leptons, and the mono-photon events, we find that the inelastic DM mass in the range of 1 MeV to 210 GeV could be tested.

**Theory / 20**

## Hybrid Renormalization for Quasi Distribution Amplitudes of A Light Baryon

**Authors:** Chao Han<sup>one</sup>; Jialu Zhang<sup>1</sup>; Wei Wang<sup>one</sup>; Yushan Su<sup>2</sup>

<sup>1</sup> 上海交通大学

<sup>2</sup> *University of Maryland*

**Corresponding Author:** elpsycongr00@sjtu.edu.cn

We develop a hybrid scheme to renormalize quasi distribution amplitudes of a light baryon on the lattice, which combines the self-renormalization and ratio scheme. By employing self-renormalization, the UV divergences and linear divergence at large spatial separations in quasi distribution amplitudes are removed without introducing extra nonperturbative effects, while making a ratio with respect to the zero-momentum matrix element can properly remove the UV divergences in small spatial separations. As a specific application, distribution amplitudes of the  $\Lambda$  baryon made of uds are investigated, and the requisite equal-time correlators, which define quasi distribution amplitudes in coordinate space, are perturbatively calculated up to the next-to-leading order in strong coupling constant  $\alpha_s$ . These perturbative equal-time correlators are used to convert lattice QCD matrix elements to the continuum space during the renormalization process. Subsequently, quasi distribution amplitudes are matched onto lightcone distribution amplitudes by integrating out hard modes and the corresponding hard kernels are derived up to next-to-leading order in  $\alpha_s$  including the hybrid counterterms. These results are valuable in the lattice-based investigation of the lightcone distribution amplitudes of a light baryon from the first principles of QCD.

**Theory / 21**

## $B$ meson anomalies and large $B^+ \rightarrow K^+ \nu \bar{\nu}$ in non-universal $U(1)'$ models

**Author:** CRISTIAN FELIPE SIERRA FONSECA<sup>1</sup>

**Co-authors:** Peter Athron<sup>1</sup>; Roberto Martinez<sup>2</sup>

<sup>1</sup> *Nanjing Normal University*

<sup>2</sup> *Universidad Nacional de Colombia*

**Corresponding Authors:** peter.athron@nju.edu.cn, remartinez@unaledu.co, cristian.sierra@nju.edu.cn

In view of both the latest LHCb measurement of  $R_{K^{(*)}}$  and the new  $2.8\sigma$  deviation reported by Belle II on  $B^+ \rightarrow K^+ \nu \bar{\nu}$  decays, we present a fit to the  $B$  meson anomalies for various one and two dimensional hypothesis including complex Wilson coefficients. We show in a model-independent way that the generic non-universal  $U(1)'$  extensions of the SM, without flavour violation, fail to simultaneously fit those observables and corroborate that they can modify  $\text{BR}(B^+ \rightarrow K^+ \nu \bar{\nu})$  up to only a 10%. In view of this deficit, we propose a new way in which those models can accommodate

the data at tree level by introducing lepton flavour violating couplings and non-diagonal elements of the charged lepton mixing matrix, with implications in future charged lepton flavour violation searches.

HF/HI/QCD / 22

## Resolving negative cross section of quarkonium hadroproduction using soft gluon factorization

**Author:** An-Ping Chen<sup>1</sup>

**Co-authors:** Yan-Qing Ma<sup>2</sup>; Ce Meng<sup>2</sup>

<sup>1</sup> Jiangxi Normal University

<sup>2</sup> Peking University

**Corresponding Author:** chenanning@jxnu.edu.cn

It was found that, using nonrelativistic QCD factorization, the predicted  $\chi_{cJ}$  hadroproduction cross section at large  $p_T$  can be negative. The negative cross sections originate from terms proportional to plus function in  ${}^3P_J^{[1]}$  channels, which are remnants of the infrared subtraction in matching the  ${}^3P_J^{[1]}$  short-distance coefficients. In this article, we find that the above terms can be factorized into the non-perturbative  ${}^3S_1^{[8]}$  soft gluon distribution function in the soft gluon factorization (SGF) framework. Therefore, the problem can be naturally resolved in SGF. With an appropriate choice of nonperturbative parameters, the SGF can indeed give positive predictions for  $\chi_{cJ}$  production rates within the whole  $p_T$  region. The production of  $\psi(2S)$  is also discussed, and there is no negative cross section problem.

Theory / 23

## Cosmological Phase Transitions in Composite Higgs Models

**Authors:** KOHEI FUJIKURA<sup>1</sup>; Yuichiro Nakai<sup>one</sup>; Ryosuke Sato<sup>2</sup>; Yaoduo Wang<sup>3</sup>

<sup>1</sup> U. Tokyo

<sup>2</sup> Osaka University

<sup>3</sup> TDLI, SJTU

**Corresponding Author:** yaoduowang@sjtu.edu.cn

We investigate cosmological phase transitions in various composite Higgs models consisting of four-dimensional asymptotically-free gauge field theories. Each model may lead to a confinementdeconfinement transition and a phase transition associated with the spontaneous breaking of a global symmetry that realizes the Standard Model Higgs field as a pseudo-Nambu-Goldstone boson. Based on the argument of universality, we discuss the order of the phase transition associated with the global symmetry breaking by studying the renormalization group flow of the corresponding linear sigma model at finite temperature, which is calculated by utilizing the  $\epsilon$ -expansion technique at the one-loop order. Our analysis indicates that some composite Higgs models accommodate phenomenologically interesting first-order phase transitions.

Theory / 24

## Theoretical Motivations for Hidden Light Bosons

**Author:** Luca Visinelli<sup>1</sup>

<sup>1</sup> *Shanghai Jiao Tong University*

**Corresponding Author:** luca.visinelli@sjtu.edu.cn

The physics beyond the Standard Model (BSM) could be represented by a hidden sector at relatively low energy scales MeV-GeV and feeble couplings to SM. Their presence in our Universe would be revealed through indirect evidences such as small oscillations of SM parameters, cosmological and astrophysical considerations, and the complementary searches in accelerators. Here, I give an overview of these BSM models including axions and hidden photons.

Plenary Session / 25

## Neutrino as a window to TeV physics: from LHC to low-energy experiments

**Author:** Gang Li<sup>1</sup>

<sup>1</sup> *Sun Yat-Sen University*

**Corresponding Author:** ligang65@mail.sysu.edu.cn

I will talk about the roles of LHC and low-energy (high-precision) experiments in searching for new physics at the TeV scale by studying the properties and interactions of neutrinos.

Theory / 26

## Dark photon effects with the kinetic and mass mixing in $Z \rightarrow \tau^- \tau^+$

**Authors:** Jin Sun<sup>1</sup>; zhi-peng Xing<sup>2</sup>

<sup>1</sup> *Shanghai Joao Tong University*

<sup>2</sup> *njnu*

**Corresponding Author:** zpxing@nmu.edu.cn

A new  $U(1)_X$  gauge boson field X can have renormalizable kinetic mixing with the standard model (SM)  $U(1)_Y$  gauge boson field Y. Besides the dark photon kinetic mixing  $\sigma$ , there could be mass mixing by introducing the additional Higgs doublet with vev engaging in  $U(1)_X$  and electroweak symmetry breaking simultaneously. The Z boson interaction with SM tau lepton is modified by defining the mixing ratio parameter  $\epsilon$ , which shows the magnitude of the mass and kinetic mixing of dark photon. We investigate the Z boson phenomenology of dark photon model with both the kinetic mixing and mass mixing. The allowed parameter region is obtained by analyzing these constraints from the vector and axial-vector couplings  $g_{V,A}^\tau$ , the decay branching ratio  $Br(Z \rightarrow \tau^- \tau^+)$  and tau lepton polarization in  $Z \rightarrow \tau^- \tau^+$ . We found that the mixing ratio plays important role in the Z boson features by choosing different  $\epsilon$  values.

Further, we attempt to find the common regions to satisfy these above four bounds for  $m_X > m_Z$  and  $m_X < m_Z$ .

However, the regions allowed by  $g_A^\tau$  and  $Br(Z \rightarrow \tau^- \tau^-)$  tends to the opposite direction so that there are not viable parameter spaces within  $2\sigma$  errors. The problem can be solved within  $3\sigma$  errors.

TeV / 27

## Probing the Higgs trilinear self-coupling through Higgs+jet production

Author: 斌周<sup>1</sup>

<sup>1</sup> 上海交通大学

Corresponding Author: 2028910235@pku.edu.cn

We present the calculation of the next-to-leading order corrections for Higgs+jet production at the Large Hadron Collider, that arise from the Higgs trilinear self-coupling ( $\lambda_{HHH}$ ).

We use the method of large top-quark mass expansion to tackle the challenging two-loop virtual amplitude, and apply the Pad\{e} approximation to extend the region of convergence of this expansion. We find that the  $\lambda_{HHH}$ -related corrections amount to 0.66% for the total cross section. For the invariant mass distribution and Higgs boson transverse momentum distribution, the corrections are mostly in the range 0.5%  $\sim$  0.7%. Our results can be used to set extra constraints on  $\lambda_{HHH}$  from the experimental data.

Theory / 28

## $B_{(s)} \rightarrow D_{(s)}^{**}$ form factors in HQEFT and model independent analysis of relevant semileptonic decays with NP effects

Author: 亚兵左<sup>1</sup>

Co-authors: 涵宇宫<sup>1</sup>; 洪瑶金<sup>1</sup>; 佳伊<sup>1</sup>; 婧赢田<sup>1</sup>; 婷婷潘<sup>1</sup>

<sup>1</sup> 辽宁师范大学

Corresponding Author: zuoyabing@lnnu.edu.cn

The form factors of  $B_{(s)}$  decays into P-wave excited charmed mesons (including  $D_0^*(2300)$ ,  $D_1(2430)$ ,  $D_1(2420)$ ,  $D_2^*(2460)$  and their strange counterparts, denoted generically as  $D_{(s)}^{**}$ ) are systematically calculated via QCD sum rules in the framework of heavy quark effective field theory (HQEFT). We consider contributions up to the next leading order of heavy quark expansion and give all the relevant form factors, including the scalar and tensor ones only relevant for possible new physics effects. The expressions for the form factors in terms of several universal wave functions are derived via heavy quark expansion. These universal functions can be evaluated through QCD sum rules. Then, the numerical results of the form factors are presented. With the form factors given here, a model independent analysis of relevant semileptonic decays  $B_{(s)} \rightarrow D_{(s)}^{**} l \bar{\nu}_l$  is performed, including the contributions from possible new physics effects. Our predictions for the differential decay widths, branching fractions and ratios of branching fractions  $R(D_{(s)}^{**})$  may be tested in more precise experiments in the future.

Theory / 29

## Muon collider signatures for a $Z'$ with a maximal $\mu - \tau$ coupling in $U(1)_{L_\mu - L_\tau}$

Author: Fei Huang<sup>1</sup>

<sup>1</sup> UJN

Theory / 30

## Precise prediction for the top quark width

Author: 烨凡王<sup>1</sup>

<sup>1</sup> Shandong university

Corresponding Author: wangyefan@sdu.edu.cn

We present the first analytic results of N<sup>3</sup>LO QCD corrections to the top-quark decay width. We focus on the dominant leading color contribution, which includes light-quark loops. At NNLO, this dominant contribution accounts for 95% of the total correction. The most precise prediction for the top-quark width is now 1.321 GeV for  $m_t = 172.69$  GeV.

Theory / 31

## Lattice study of singlet-assisted electroweak phase transition

Authors: Guotao Xia<sup>1</sup>; Lauri Niemi<sup>one</sup>; Michael J. Ramsey-Musolf<sup>one</sup>

<sup>1</sup> TDLI, SJTU

Corresponding Author: xiagt-summer@sjtu.edu.cn

In this study, we use lattice to reveal nonperturbative information of the electroweak phase transition in the real-singlet extension of the Standard Model, based on the 2-loop 3D EFT framework.

Importantly, the new information is that the lattice determines the true nature of the electroweak phase transition, capable to identify it as the first order type or not, an important qualitative behavior to which perturbation theory is blind. In scenarios where perturbation theory implies a weakly first order phase transition, lattice is always more reliable than the perturbation theory. In this regime, the symmetry-breaking transition may be crossover rather than a true phase transition. On the other hand, for strong transitions, both methods yield quantitatively close results, particularly when 2-loop perturbation theory is used.

This nonperturbative framework holds potential for other Higgs-sector extensions of the SM. Besides, by holding two powerful tools, 2-loop perturbation scanning and lattice, we will explore associated phenomenology in the future.

Theory / 32

## Observation of new structure in the $J/\psi$ / $\psi$ mass spectrum in proton-proton collisions at $\sqrt{s} = 13$ TeV

Author: Xining Wang<sup>1</sup>

<sup>1</sup> Tsinghua University

Corresponding Author: xining.wang@cern.ch

A search is reported for near-threshold structures in the  $J/\psi$ / $\psi$  invariant mass spectrum produced in proton-proton collisions at  $\sqrt{s} = 13$  TeV from data collected by the CMS experiment, corresponding to an integrated luminosity of 135 fb<sup>-1</sup>. A new structure is observed with a significance above 5 standard deviations at a mass of  $6552 \pm 10$  (stat)  $\pm 12$  (syst) MeV. Another structure with even higher significance is found at a mass of  $6927 \pm 9$  (stat)  $\pm 4$  (syst) MeV, which is consistent with the X(6900



) resonance reported by the LHCb experiment and confirmed by the ATLAS experiment. Evidence for another new structure, with a local significance of 4.1 standard deviations, is found at a mass of  $7287+20 -18$  (stat)  $\pm 5$  (syst) MeV. The masses and significances are obtained in a model without considering possible quantum mechanical interference between the resonances. Incorporating this interference provides a better description of the mass spectrum between the resonances and shifts the measured masses by up to 150 MeV.

Theory / 33

## Electroweak sphalerons, scalar multiplets, and symmetry breaking patterns

**Authors:** Yanda Wu<sup>1</sup>; Wenxing Zhang<sup>2</sup>; Michael Ramsey-Musolf<sup>2</sup>

<sup>1</sup> TDLI, SJTU

<sup>2</sup> TDLI/SJTU

**Corresponding Author:** yanda.wu7@sjtu.edu.cn

In this study, we present a comprehensive analysis of the electroweak sphaleron formalism and its application to electroweak phase transition (EWPT) patterns in extensions of the Standard Model scalar sector with electroweak multiplets. We offer an equivalence proof for different choices for the form of sphaleron configurations; construct the previously unestablished high-dimensional SU(2) sphaleron transformation matrix; investigate the scalar multiplet topology map and baryon number charge relation; and revisit the required boundary conditions needed for solving the sphaleron field equations. We then scrutinize the leading order sphaleron dynamics in the context of a multi-step EWPT. We showcase two distinct analytical approaches for extending the SU(2) scalar multiplet to the standard model (SM) under differing EWPT scenarios, and perform an explicit calculation of the sphaleron energy using a septuplet example. In the context of a single-step EWPT leading to a mixed phase, we find that the additional multiplet's contribution to the sphaleron energy is negligible, primarily due to the prevailing constraint imposed by the  $\rho$  parameter. Conversely, in a two-step EWPT scenario, the sphaleron energy can achieve significantly high values during the initial phase, thereby markedly preserving baryon asymmetry if the universe undergoes a first-order EWPT. In both cases, we delineate the relationship between the sphaleron energy and the parameters relevant to dark matter phenomenology.

TeV / 34

## Search for Higgs boson pair production in the $b\bar{b}\mu\mu$ final state at the LHC (Remote)

**Author:** Botao Guo<sup>1</sup>

<sup>1</sup> Peking University

**Corresponding Author:** botao.guo@stu.pku.edu.cn

The Higgs boson pair production via gluon-gluon fusion and vector boson fusion in the  $b\bar{b}\mu\mu$  final state at the LHC is studied to probe the Higgs self-coupling  $\kappa\lambda$  and the four-boson HHVV coupling  $\kappa_2V$  for the first time. A cut-based analysis and a machine-learning analysis using boosted decision trees are performed with categorizations and optimizations depending on the variations of these couplings. The expected sensitivities are extracted with different integrated luminosities assumed up to the full highluminosity LHC runs. The expected upper limit at 95% confidence level on the Higgs boson pair production is calculated as 47 (28) times the Standard Model cross section using the cut-based method (boosted decision trees) for the gluon-gluon fusion production and 928 for the vector boson fusion production, assuming an integrated luminosity of  $3000 \text{ fb}^{-1}$ . The expected

constraints on the couplings at 95% confidence level are calculated to be  $-13.8 < \kappa\lambda < 19.1$  ( $-10.0 < \kappa\lambda < 15.5$ ) and  $-3.4 < \kappa 2V < 5.5$  using the cut-based method (boosted decision trees), respectively, assuming an integrated luminosity of 3000 fb<sup>-1</sup>.

Theory / 35

## MatchingEFT.jl: An Automated Tool for Tree and One-Loop Level Matching

Author: 成杰杨<sup>1</sup>

<sup>1</sup> 高能物理研究所 & 理论物理研究所

Corresponding Author: yangchengjie@ihep.ac.cn

We present MatchingEFT.jl, an automated tool to extract hard region contribution of the tree-level and one-loop 1PI amplitude, which can be matched to the standard model effective theory (SMEFT) operator basis in ABC4EFT.

FeAnGen4EFT performs the feynman diagram generation for the specific scattering process using the designed universal feynrules output file (UFO) and Qgraf. It offers the designated Feynman diagrams' amplitudes explicitly to provide non-trivial check for the gauge invariance. It will use FORM script to process extracting hard region expansion for the output from the previous step. These results will be given in a physical basis such as P-basis and Y-basis in ABC4EFT by using the on-shell amplitude basis method for a further crosscheck. FeAmGen4SMEFT has been built with lightweight, generality, flexibility, specialization, and efficiency in mind.

These ingredients allow FeAmGen4SMEFT to have more applications beyond the matching based on the complete operator basis with the specific mass dimension offered by ABC4EFT. One of these applications includes the one-loop renormalization of arbitrary theories.

We have performed one-loop matching for some processes under the specified UV model and obtained consistent results with other papers.

Theory / 36

## Elliptic anisotropy of hard probes from parton scatterings in small collision systems

Author: 思宇汤<sup>1</sup>

<sup>1</sup> 武汉纺织大学

Corresponding Author: tsy@wtu.edu.cn

The hard probes, including jets and heavy flavors, play an important role in investigating the properties of quark-gluon plasma (QGP) formed in heavy-ion collisions. The positive elliptic flow of hard probes observed in semi-central Pb-Pb collisions indicates that the hard partons suffered strong interactions in the deconfined QCD medium and then obtain the collectivity. However, recent measurements show also a non-zero  $v_2$  for high- $p_T$  charged particles and heavy flavor hadrons in high-multiplicity p-Pb collisions for both mid and forward rapidities, whose origin is still debated.

In this contribution, we employ a multi-phase transport model (AMPT) to calculate the  $v_2$  of jet particles and open heavy-flavour hadron decay muons in p-Pb collisions at mid and forward rapidity, respectively. The results are obtained using the two-particle correlation method and the advanced nonflow subtraction strategy. We will systematically introduce how the collectivity of hard partons are generated from parton scatterings, and then propagated to the final state in small collision systems. Comparisons with experimental results will be presented as well. This work will provide further insights into understanding the origin of elliptic anisotropy of hard probes in small collision systems, and has referential value for the future measurements.

Poster Session / 37

## Improved Asymptotic Formulae for Statistical Interpretation Based on Likelihood Ratio Tests

Authors: LIGANG XIA<sup>1</sup>; yan zhang<sup>2</sup>

<sup>1</sup> Nanjing University

<sup>2</sup> 南京大学

Corresponding Author: yanz@cern.ch

In this work, we try to improve the classic asymptotic formulae to describe the probability distribution of likelihood-ratio statistical tests. The idea is to split the probability distribution function into two parts. One part is universal and described by the asymptotic formulae. The other part is case-dependent and estimated explicitly using a 6-bin model proposed in this work. The latter is similar to doing toy simulations and hence is able to predict the discrete structures in the probability distributions. The new asymptotic formulae provide a much better differential description of the test statistics. The better performance is confirmed in two toy examples.

Theory / 38

## Limiting FCNC induced by a CP symmetry of order 4

Author: Duanyang Zhao<sup>1</sup>

<sup>1</sup> 中山大学

Corresponding Author: zhaody8@mail2.sysu.edu.cn

CP4 3HDM is a three-Higgs-doublet model based on the CP symmetry of order 4 (CP4). Imposing CP4 leads to remarkable connections between the scalar and Yukawa sectors and unavoidably generates tree-level flavor-changing neutral couplings (FCNC). It remains unclear whether FCNC can be sufficiently suppressed in the CP4 3HDM. In this paper, we systematically explore this issue. We first develop an efficient scanning procedure which takes the quark masses and mixing as input and expresses the FCNC matrices in terms of physical quark observables and quark rotation parameters. This procedure allows us to explore the FCNC effects for all the Yukawa sectors possible within the CP4 3HDM. We find that, out of the eight possible CP4 Yukawa sectors, only two scenarios are compatible with the K, B, Bs and, in particular, D-meson oscillation constraints. The results of this work serve as clear guidelines for future phenomenological scans of the model.

Theory / 39

## Aiming for Tops of ALPs with a Muon Collider

Authors: So Chigusa<sup>one</sup>; Sudhakantha Girmohanta<sup>1</sup>; Yufei Zhang<sup>one</sup>; Yuichiro Nakai<sup>one</sup>

<sup>1</sup> Tsung-Dao Lee Institute and Shanghai Jiao Tong University

Corresponding Author: yufei.zhang@sjtu.edu.cn

Future muon colliders with center-of-mass energy of  $\mathcal{O}(1-10)$  TeV can provide a clean high-energy environment with advantages in searches for TeV-scale axion-like particles (ALPs), pseudo-Nambu-Goldstone bosons associated with spontaneously broken global symmetries, which are widely predicted in physics beyond the Standard Model (SM).

We exploit ALP couplings to SM fermions, and guided by unitarity constraints, build a search strategy focusing on the ALP decay to top quark pairs.

Theory / 40

## The installation progress of upgrade Upstream Tracker at LHCb Upgrade II

Author: Xiaojie Jiang<sup>1</sup>

<sup>1</sup> IHEP(Beijing)

Corresponding Author: jiangxj@ihep.ac.cn

The Upstream Tracker is a key component of the LHCb updating. Based on silicon strip sensors, the UT will contribute to efficient and high-speed track reconstruction. UT's installation is a long-term and challenging work. Fortunately, the installation of UT is completed earlier this year. It also passed many kinds of electronics and service tests. The details of installation and pre-tests for the UT detector will be introduced.

Theory / 41

## Explaining the CDF $W$ -mass shift and $(g-2)_\mu$ in a $Z'$ scenario and its implications for the $b \rightarrow s \ell^+ \ell^-$ processes

Author: 袁兴博<sup>1</sup>

<sup>1</sup> CCNU

Corresponding Author: y@ccnu.edu.cn

In the past few years, several indirect hints for New Physics beyond the SM arose in precision measurements, e.g.,  $(g-2)_\mu$  and the  $W$ -boson mass. In this work, we consider a model containing new vector-like Fermion partner gauged under a new  $U(1)'$  symmetry. It is found that the latest CDF  $m_W$  measurement and  $(g-2)_\mu$  can be simultaneously accommodated. We have also considered several other experimental constraints, including the neutrino trident production,  $Z \rightarrow \mu\mu$  decay, dimuon resonance searches at the LHC, etc. Implications for the  $b \rightarrow s \ell^+ \ell^-$  process will be discussed. (This work is based on 2205.02205 and 2307.05290.)

Theory / 42

## Massive Scattering Amplitudes for Standard Model: On-shell Massless-Massive Correspondence

Authors: 江浩于<sup>1</sup>; 超鄂<sup>1</sup>; 郁涵倪<sup>1</sup>

<sup>1</sup> Institute of Theoretical Physics, Chinese Academy of Sciences

Corresponding Author: niyuhan@itp.ac.cn

We organize massive tree-level amplitudes in Standard Model by power counting and helicity category, and match them with their high energy origins. The construction of massive amplitudes

is based on the massive bootstrap method, decomposing internal and external structures, and the existing leading orders of massless-massive correspondence. For the matching of higher order components, we introduce the on-shell Higgsing mechanism first proposed by (R. Balkin et al., 2022) and further the discussion to propagators.

Theory / 43

## The Prototype Design of PEB - a Component of the HGTD In-detector Electronics for the ATLAS Phase-II Upgrade

Author: Zhenwu Ge<sup>1</sup>

<sup>1</sup> Nanjing University

Corresponding Author: 602022220019@smail.nju.edu.cn

The High Granularity Timing Detector (HGTD) is proposed as a part of the ATLAS Phase-II upgrade to mitigate the impact of pileup on object reconstruction by precisely measuring the time of tracks. In addition, HGTD also provides an instantaneous measurement of the luminosity. HGTD is composed of 8032 front-end modules. Each module consists of two Low Gain Avalanche Detectors (LGADs) of approximately  $2 \times 2 \text{ cm}^2$  bump-bonded to two ATLAS LGAD Timing Integrated Read-Out Chips (ALTIROC) and held together by a module flex (flexible PCB). Each module will be connected to the Peripheral Electronics Boards (PEB) through a flex tail (another flexible PCB). The connections between on-detector and off-detector electronics are performed via optical fibers, high/low voltage cables, interlock cables and monitoring signal cables. The PEB acts as a bridge between the front-end modules and the off-detector systems. The optical fibers provide shared data streams for Timing, Trigger and Control (TTC), Detector Control System (DCS) and Data Acquisition System (DAQ), and dedicated data streams for the luminosity system. The PEB uses the low-power GigaBit Transmission chip (lpGBT) and the Versatile Link + Transceiver (VTRx+). The PEB also includes the 12 V to 1.2 V DC-DC converters (bPOL12v) for the digital and analogue voltages supplied to the front-end modules. The supply voltages are monitored using the internal multiplexed ADC on the lpGBTs. Since the input channel number of this ADC is limited to 8, a multiplexing chip is required to handle all the signals connected to PEB. A full custom 64-to-1 multiplexing ASIC (MUX64) has been developed with a radiation tolerance suitable for its implementation on the PEB. According to the optimization of mirror structure for the layout of the modules, 6 types of PEBs need to be designed for HGTD. Based on previous development experience, the PEB 1F was chosen to be designed first as a prototype since it is the most complicated PEB type, which supports up to 55 front-end modules with 12 lpGBT, 9 VTRx+ and 52 bPOL12v in a very limited space. The requirements and overall specifications of the electronics of HGTD will be presented as well as the technical design and the project status.

Theory / 44

## Dark matter distribution, structure formation and the potential to distinguish thermal histories of dark matter.

Authors: Fei Huang<sup>1</sup>; Haolin Li<sup>2</sup>; Yong Du<sup>3</sup>; Yuan-Zhen Li<sup>one</sup>; 江浩于<sup>4</sup>

<sup>1</sup> Weizmann Institute of Science

<sup>2</sup> ITP-CAS

<sup>3</sup> TDLI-SJTU

<sup>4</sup> Institute of Theoretical Physics, Chinese Academy of Sciences

Corresponding Author: liyuanzhen@itp.ac.cn

It is important to understand the implications of current observational constraints and potential signatures on the thermal history of dark matter. Using the freeze-in/-out scenarios as templates, we

revisit dark-matter production by solving the Boltzmann equations at the level of the phase-space distribution. We also investigate the current Lyman-alpha constraints on mass of the dark matter and build the connection between the mass and the production mechanism of dark matter and find that the current observation on structure formation can be imposed to constrain the decoupling temperatures and the phase-space distribution of dark matter. We further explore the potential of distinguishing different possible thermal histories of dark matter with hypothetical future observational data. This method can be more generally applied to other scenarios.

**Theory / 46**

## Double Parton Scattering Effect in the Measurement of W-Mass

**Author:** Rui Zhang<sup>1</sup>

**Co-author:** Hao Zhang<sup>2</sup>

<sup>1</sup> IHEP

<sup>2</sup> Theoretical Physics Division, Institute of High Energy Physics, Chinese Academy of Sciences

**Corresponding Author:** zhangr@ihep.ac.cn

Recently, the  $W$  boson mass measured by the CDF-II collaboration shows large tension with the standard model prediction and other measurements. In this work, we look into the double parton scattering (DPS) contribution in CDF-II  $W$  mass measurement. We show that the DPS process can increase the measured mass as  $\Delta M_W = 20 - 200$  MeV for the missing transverse momentum fit and  $\Delta M_W = 0 - 50$  MeV for the transverse mass fit. It is comparable to the  $W$ -mass tension and should be take into consideration. The DPS effect can also appears in other inclusive measurements, since it contributes  $\sim 10^{-2}$  events in total and cause a  $\mathcal{O}(10^{-2}) - \mathcal{O}(10^{-1})$  GeV shift of the missing transverse momentum.

**Theory / 47**

## Hunting for sterile neutrino at colliders

**Author:** Cong-Feng QIAO<sup>1</sup>

<sup>1</sup> UCAS

**Corresponding Author:** qiaocf@ucas.ac.cn

We study the feasibility of observing sterile neutrino at the high energy colliders, using direct and indirect production channels in heavy meson/baryon and Higgs decays. It is found that these processes may set certain new constraints on the mass of sterile neutrino in present running and next generation experiments.

**Theory / 48**

## Probing Neutral Triple Gauge Couplings at the LHC, CEPC and SPCC

**Author:** Ruiqing Xiao<sup>one</sup>

**Corresponding Author:** 343673681@qq.com

We study probes of neutral triple gauge couplings (nTGCs) at the LHC, CEPC and SPPC. The nTGCs provide a unique window to the new physics beyond the Standard Model (SM) because they can arise from SM effective field theory (SMEFT) operators that respect the full electroweak gauge group  $SU(2)_L \otimes U(1)_Y$  of the SM only at the level of dimension-8 or higher. We derive the neutral triple gauge vertices (nTGVs) generated by these dimension-8 operators in the broken phase and map them onto a newly generalized form factor formulation, which takes into account only the residual  $U(1)_{em}$  gauge symmetry. Using this mapping, we derive new relations between the form factors that guarantee a truly consistent form factor formulation of the nTGVs and remove large unphysical energy-dependent terms. We then analyze the sensitivity reaches of the LHC, CEPC and SPPC for probing the nTGCs via both the dimension-8 nTGC operators and the corresponding nTGC form factors. We compare their sensitivities with the existing LHC measurements of nTGCs and with those of future colliders.

**Theory / 50**

## **$B \rightarrow D$ form factors beyond leading power and extraction of $|V_{cb}|$ and $R(D)$**

**Corresponding Author:** yanbing.wei@bjut.edu.cn

**Theory / 51**

## **Unveiling time-varying signals of ultralight bosonic dark matter at collider and beam dump experiments**

**Theory / 52**

## **Heavy long-lived coannihilation partner from inelastic Dark Matter model and its signatures at the LHC**

**Theory / 53**

## **利用有限温度有效理论研究电弱相变**

**Corresponding Author:** 490555870@qq.com

**Theory / 54**

## **BBN 和 CMB 限制低能标新物理**

**Corresponding Author:** 2080226955@qq.com

Theory / 55

## **Testing Complex Singlet Scalar Cosmology at the Large Hadron Collider**

Corresponding Author: zhangwenxing@sjtu.edu.cn

Theory / 56

## **Massive gauge theory with quasigluon for hot SU( N ): Phase transition and thermodynamics**

Corresponding Author: jackpotzhujiang@gmail.com

Theory / 57

## **Electroweak phase transitions proceed via bubbles vs. domain walls (Remote)**

Corresponding Author: jiangyun5@sysu.edu.cn

Plenary Session / 58

## **Theory overview: Collider Physics**

Corresponding Author: yanbin@ihep.ac.cn

Plenary Session / 59

## **Precision Higgs measurements at the LHC**

Corresponding Author: xiaohu.sun@cern.ch

Plenary Session / 60

## **Search for HH production at the LHC**

Corresponding Author: nan.lu@cern.ch

Plenary Session / 61



## **BSM/rare Higgs**

**Corresponding Author:** yanlin@cern.ch

Plenary Session / 62

## **Theory overview: Heavy flavor physics**

**Corresponding Author:** shenylmeteor@ouc.edu.cn

Plenary Session / 63

## **重味谱学**

**Corresponding Author:** liupan.an@cern.ch

Plenary Session / 64

## **Search for rare decays at LHCb**

**Corresponding Author:** jibo.he@ucas.ac.cn

Plenary Session / 65

## **ALICE 实验中集体运动研究进展**

**Corresponding Author:** shouqiye@fudan.edu.cn

Plenary Session / 66

## **Theory overview: QCD**

**Corresponding Author:** haitao.li@sdu.edu.cn

Precision prediction for top quark physics

Top quark is the heaviest particle in the standard model. Once produced, the top quark immediately decays to a W boson and a bottom quark before hadronization. Precision studies for top quark physics plays essential role in the validity of the Standard Model and exploring potential avenues for new physics. In this talk I will review recently theoretical development in top quark physics, which includes the theoretical efforts for top quark production and decay.

**Plenary Session / 67**

## **SM measurements at the LHC (+Top)**

**Corresponding Author:** chen.zhou@cern.ch

**Plenary Session / 68**

## **CPV**

**Corresponding Author:** wenbin.qian@ucas.ac.cn

**Plenary Session / 69**

## **CEPC review**

**Plenary Session / 70**

## **ATLAS upgrade**

**Corresponding Author:** lailin.xu@cern.ch

**Plenary Session / 71**

## **CMS upgrade**

**Corresponding Author:** wangfeng@ihep.ac.cn

**Plenary Session / 72**

## **LHCb upgrade**

**Corresponding Author:** yangzw@pku.edu.cn

**Plenary Session / 73**

## **ALICE upgrade**

**Corresponding Author:** xiao\_mei\_li@foxmail.com

Plenary Session / 74

## **Theory overview: Heavy ion physics**

Corresponding Author: weiyaoke@ccnu.edu.cn

Plenary Session / 75

## **ALICE 实验中奇异强子的产生**

Corresponding Author: zbyin@mail.ccnu.edu.cn

Plenary Session / 76

## **ALICE 实验中重夸克的产生与输运**

Corresponding Author: xinye.peng@cern.ch

Plenary Session / 77

## **Heavy Ion physics at ATLAS and CMS**

Corresponding Author: qipenghu@ustc.edu.cn

Plenary Session / 78

## **Theory Overview: Neutrino Physics**

Corresponding Author: ligangpku@163.com

Plenary Session / 79

## **SUSY results at LHC**

Corresponding Author: xuda@ihep.ac.cn

Plenary Session / 80

## **Dark matter and unconventional searches at the LHC**

Corresponding Author: nzhou@sjtu.edu.cn

**Plenary Session / 81**

## **Exotics (non-SUSY) searches at the LHC (+Flavor)**

**Corresponding Author:** shencp@fudan.edu.cn

**Plenary Session / 82**

## **Speech from SJTU-TDLI and CPS**

**Corresponding Author:** jzhang1@sjtu.edu.cn

**Plenary Session / 83**

## **Speech from Funding Agency**

**Plenary Session / 84**

## **Speech from CPS High Energy Branch**

**Plenary Session / 85**

## **Announcements from LOC**

**Corresponding Authors:** haijun.yang@sjtu.edu.cn, shuli@sjtu.edu.cn

**Plenary Session / 86**

## **ATLAS report**

**Corresponding Author:** wuyusheng@ustc.edu.cn

**Plenary Session / 87**

## **CMS report**

**Corresponding Author:** kai.yi.yik@gmail.com

**Plenary Session / 88**

## **LHCb report**

**Corresponding Author:** hengne.li@m.scnu.edu.cn

**Plenary Session / 89**

## **ALICE report**

**Corresponding Author:** song\_zhang@fudan.edu.cn

**Plenary Session / 90**

## **Theory overview: New Physics at the LHC: SMEFT and Beyond**

**Corresponding Author:** hjhe@sjtu.edu.cn

**Plenary Session / 91**

## **Machine Learning at LHC**

**Corresponding Author:** rui.zhang@cern.ch

**Closing / 92**

## **CLHCP2023 Summary**

**Corresponding Author:** yuanning.gao@pku.edu.cn

**Closing / 93**

## **Announcements**

**Corresponding Authors:** jiali@pku.edu.cn, zhanghq@mail.ihep.ac.cn

**Instrumental / 94**

## **Status of USTC-IME Pre-production sensor for the ATLAS High Granularity Timing Detector**

**Corresponding Author:** kuo.ma@cern.ch

**Instrumental / 95**

## **First results of USTC AC-coupled LGADs**

**Instrumental / 96**

## **Electronics test results of Altiroc2 and Altiroc3**

**Instrumental / 97**

## **HGTD module assembly and module test at USTC**

**Corresponding Author:** aonan.wang@cern.ch

**Instrumental / 98**

## **ATLAS High granularity timing detector activity at IHEP/NJU**

**Corresponding Author:** liangzj@ihep.ac.cn

**Instrumental / 99**

## **IHEP-IME radiation hard LGAD sensor pre-production for ATLAS High granularity timing detector**

**Instrumental / 100**

## **The Prototype Design of PEB - a Component of the HGTD In-detector Electronics for the ATLAS Phase-II Upgrade**

**Corresponding Author:** 602022220019@smail.nju.edu.cn

**Instrumental / 101**

## **R&D of AC-LGAD based 4D tracker with precision timing information for HL-LHC and the future colliders**

**Corresponding Author:** sunwy@ihep.ac.cn

**Instrumental / 102**

## **Module performance study for the High Granularity Timing Detector of ATLAS**

**Corresponding Author:** yangxuan@ihep.ac.cn

**Instrumental / 103**

## **Radiation Tolerance of the MUX64 for the High Granularity Timing Detector of ATLAS**

**Corresponding Author:** chuanye.wang@cern.ch

**Instrumental / 104**

## **Automatic Module assembly development for ATLAS High granularity timing detector**

**Corresponding Author:** xinhui.huang@cern.ch

**Instrumental / 105**

## **HGTD PEB DC/DC Power Block in Low Temperature and Magnetic Field Operation**

**Corresponding Author:** zhaimingjie@ihep.ac.cn

**Instrumental / 106**

## **Performance of USTC-IME sensors in test beams at DESY and CERN**

**Corresponding Author:** zxx0209@mail.ustc.edu.cn

**Instrumental / 107**

## **CMS Endcap Timing Layer of MTD**

**Corresponding Author:** yuchengjun@ustc.edu.cn

**Instrumental / 108**

## **MTD sensor and assembly**

**Corresponding Author:** jinwang07@pku.edu.cn

**Instrumental / 109**

## **HGCal MAC status at IHEP**

**Corresponding Author:** yutz@ihep.ac.cn

**Instrumental / 110**

## **HGCal SQC at IHEP and domestic silicon sensor R&D for calorimeter**

**Corresponding Author:** 202183900031@sdu.edu.cn

**Instrumental / 111**

## **Computing vision QC in HGCal production**

**Corresponding Author:** xhyuan@ihep.ac.cn

**Instrumental / 112**

## **基于 SystemC 对 FOCAL-E 像素层进行模拟分析**

**Corresponding Author:** jie.yi@cern.ch

**Instrumental / 113**

## **FOCAL 探测器像素层数据分析现状**

**Corresponding Author:** liuhe@cern.ch

**Instrumental / 114**

## **ECAL simulation for LHCb Upgrade II**

**Corresponding Author:** jfei@cern.ch



**Instrumental / 115**

## **Testbeam results of the LHCb Upgrade II ECAL prototype**

**Corresponding Author:** ge.ma@cern.ch

**Instrumental / 116**

## **Performance of the GAGG crystal for LHCb Upgrade II ECAL**

**Corresponding Author:** yuanzy@pku.edu.cn

**Instrumental / 117**

## **ATLAS ITk Strip Module Site Qualification at IHEP**

**Corresponding Author:** guolei@ihep.ac.cn

**Instrumental / 118**

## **Cold Noise Study in ATLAS ITk Strip Detectors**

**Corresponding Author:** lizhan@ihep.ac.cn

**Instrumental / 119**

## **SEE test of production ASICs in CSNS**

**Corresponding Author:** psg19@mails.tsinghua.edu.cn

**Instrumental / 120**

## **A study on the feasibility of CSNS becoming an ATLAS ITk sensor QA irradiation site**

**Corresponding Author:** hui-li20@mails.tsinghua.edu.cn

**Instrumental / 121**

## **Installation of LHCb Upstream Tracker**

**Corresponding Author:** jiangxj@ihep.ac.cn

**Instrumental / 122**

## **Simulation of UT for LHCb Upgrade II**

**Corresponding Author:** shengshuqi@ihep.ac.cn

**TeV / 124**

## **Early Run3 $H \rightarrow \gamma\gamma$ fiducial Cross-Section measurement**

**TeV / 125**

## **Test of CP Invariance in Higgs Boson Vector-Boson-Fusion Production Using $\gamma\gamma$ Channel with the ATLAS Detector**

**TeV / 126**

## **Novel photon energy calibration method and Higgs mass measurement**

**Corresponding Author:** hemx@ihep.ac.cn

**TeV / 127**

## **(Remote) Measurement of the $H \rightarrow ZZ \rightarrow 4l$ cross-sections in pp collisions at 13 TeV**

**TeV / 128**

## **Measurements of inclusive and differential cross sections for the Higgs boson production and decay to four-leptons in proton-proton collisions at $\sqrt{s} = 13$ TeV**

**TeV / 129**

**Legacy search for the non-resonant production of Higgs boson pairs via gluon fusion bosonfusion in the  $\tau\tau\tau\tau$  final state in proton-proton collisions at  $\sqrt{s} = 13$  TeV with the ATLAS detector (remote)**

Corresponding Author: lihan@cern.ch

TeV / 130

**Studies of new Higgs boson interactions through nonresonant HH production in final state in pp collisions at  $\sqrt{s} = 13$  TeV with the ATLAS detector**

Corresponding Author: zihang.jia@cern.ch

TeV / 131

**Preliminary Results on Higgs Pair Production in Multi-lepton Channel with the ATLAS Experiment**

Corresponding Author: zc\_1994@163.com

TeV / 132

**Search for diHiggs with VHH (Remote)**

Corresponding Author: licheng.zhang@cern.ch

TeV / 133

**Higgs mass and width measurement in ZZ to 4-leptons final state with full Run2 data (CMS)**

Corresponding Author: chenguang.zhang@ihep.ac.cn

TeV / 134

**Measurement of Higgs boson mass and width with LHC run2 data at the ATLAS e**

Corresponding Author: zhangyf42@mail.ustc.edu.cn

HF/HI/QCD / 135

## **$B(s) \rightarrow D(^*)\phi$**

HF/HI/QCD / 136

## **Charm spectroscopy at LHCb**

HF/HI/QCD / 137

## **Interference fit to the double $J\psi$ mass spectrum at CMS**

Corresponding Author: xining.wang@cern.ch

HF/HI/QCD / 138

## **b-hadron FCNC decays at LHCb**

HF/HI/QCD / 139

## **$D_s$ and $D^+$ production in pPb**

Corresponding Author: yiluo@cern.ch

HF/HI/QCD / 140

## **$\Psi(2S)/J\psi$ versus multiplicity in pp**

Corresponding Author: youen.kang@cern.ch

HF/HI/QCD / 141

## **$B_s \rightarrow \phi\phi$**

Corresponding Author: kechen.li@cern.ch

HF/HI/QCD / 142

## **lhcb EW results**

HF/HI/QCD / 143

## **Associated quarkonium production**

Corresponding Author: jialu.wang@cern.ch

HF/HI/QCD / 144

## **Dicharm decay of beauty baryons**

Corresponding Author: yiduo.shang@cern.ch

HF/HI/QCD / 145

## **Omegac0 two-body hadronic decays at LHCb**

Corresponding Author: chuangxin.lin@cern.ch

HF/HI/QCD / 147

## **$B \rightarrow D^* K$**

HF/HI/QCD / 148

## **$B_s \rightarrow D_s 1K$**

Corresponding Author: zirui.w@cern.ch

Remote

TeV / 149

## **Search for the exotic decay of the Higgs boson into a Z boson and a light pseudo-scalar decaying into two photons in pp collisions at 13TeV (CMS)**

Corresponding Author: wangzebing@ihep.ac.cn

TeV / 150

## **Search for extra Higgs bosons through same-sign top-quark production in association with an extra jet(Remote)**

Corresponding Author: meng.lu@cern.ch

TeV / 151

## **Searches for lepton flavour violation in Higgs boson decays, $H \rightarrow e\tau$ and $H$**

Corresponding Author: antonio.de.maria@cern.ch

TeV / 152

## **Searches for Lepton-flavour-violating decays of the Higgs boson**

Corresponding Author: minlin.wu@cern.ch

TeV / 153

## **Search for a standard model-like Higgs boson in the mass range between 70 and 110 GeV in the diphoton final state in proton-proton collisions at 13 TeV (CMS)**

Corresponding Author: taojq@ihep.ac.cn

TeV / 154

## **Evidence for the Higgs boson decay to a Z boson and a photon at the LHC**

Corresponding Author: hemx@ihep.ac.cn

TeV / 156

## **Run2 combination of the Higgs boson decay to a Z boson and a photon at the LHC**

Corresponding Author: mtzhang@stu.pku.edu.cn

TeV / 157

## **Interpretations of the measurements of Higgs boson production and decay rates and differential cross-sections based on the Nature paper(remote)**

Corresponding Author: zyjonah@sjtu.edu.cn

TeV / 158

## **CMS Higgs and double Higgs combinations (Remote)**

Corresponding Author: jialin.guo@ihep.ac.cn

TeV / 159

## **Measurement of ttW/ttH production in multilepton final states with ATLAS Run 2 data**

TeV / 160

## **Measurement of Higgs boson decaying into $b\bar{b}$ using Run2 dataset at the ATLAS experiment**

HF/HI/QCD / 161

## **Non-prompt $\Lambda_c^+$ production with machine learning in p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV with ALICE**

Corresponding Author: lupz@mail.ustc.edu.cn

HF/HI/QCD / 162

## **ALICE 实验超氙产生测量 (remote)**

Corresponding Author: 20210200009@fudan.edu.cn

HF/HI/QCD / 163

## **Study of heavy-flavor physics via semi-muonic decays with ALICE**

**Corresponding Author:** maolin.zhang@cern.ch

HF/HI/QCD / 164

### **ALICE 实验轻核动量关联研究 (Remote)**

**Corresponding Author:** dfwang20@fudan.edu.cn

HF/HI/QCD / 165

### **Measurement of $\Omega_c$ branching-fraction ratio at ALICE**

**Corresponding Author:** chengtiantian@mails.cnu.edu.cn

HF/HI/QCD / 166

### **Pinning down the origin of collectivity in small systems with ALICE(Remote)**

**Corresponding Author:** mingrui.zhao@cern.ch

HF/HI/QCD / 167

### **Probing the nuclear structure with multiparticle correlation in Xe–Xe collisions**

**Corresponding Author:** liscklu@hotmail.com

HF/HI/QCD / 168

### **基于 run3 数据的强子-奇异粒子关联研究现状**

**Corresponding Author:** kaicui@mails.cnu.edu.cn

HF/HI/QCD / 169

### **Measurement of inclusive $J/\Psi$ and $\Psi(2S)$ production at midrapidity in pp collisions at 13.6 TeV with ALICE**

**Corresponding Author:** zhang\_yuan@mail.ustc.edu.cn



HF/HI/QCD / 170

## **Non-prompt $J/\Psi$ production in pp collisions with ALICE**

Corresponding Author: wenda.guo@uib.no

HF/HI/QCD / 171

## **Measurement of inclusive $J/\Psi$ polarization at midrapidity in pp collisions at 13.6 TeV with ALICE**

Corresponding Author: zhenjun.xiong@cern.ch

HF/HI/QCD / 172

## **$\Xi_c^0$ production vs multiplicity via hadronic decay in pp at 13 TeV**

Corresponding Author: tao.fang@cern.ch

HF/HI/QCD / 173

## **Study of strange hadron production in jets and the underlying events with ALICE**

Corresponding Author: langxu@mails.ccnu.edu.cn

HF/HI/QCD / 174

## **ALICE 实验小系统碰撞中集体流研究 (Remote)**

Corresponding Author: 19110200003@fudan.edu.cn

HF/HI/QCD / 175

## **ALICE 实验手征反常效应研究 (Remote)**

Corresponding Author: wangcz22@m.fudan.edu.cn

HF/HI/QCD / 176

## **Measurement of the prompt and non-prompt $J/\Psi$ production in Pb–Pb collisions at 5.02 TeV with ALICE**

Corresponding Author: zhusenjie@mail.ustc.edu.cn

HF/HI/QCD / 177

## **Study of jet quenching effects using jet-hadron correlations with ALICE(Remote)**

Corresponding Author: yongzhen.hou@cern.ch

HF/HI/QCD / 178

## **Study of beauty quark production properties via non-prompt D mesons with ALICE**

Corresponding Author: jon.harris@mails.ccnu.edu.cn

Theory / 181

## **New perspectives on UV divergences of loops and the hierarchy problem**

Corresponding Author: jialb@mail.nankai.edu.cn

TeV / 182

## **[Cancel] Mono-X Signatures of an Absorbed Fermionic Dark Matter at the LHC**

Corresponding Author: makainca@yeah.net

TeV / 183

## **Search for direct production of electroweakinos in final states with one lepton, jets and missing transverse momentum with the ATLAS detector**

Corresponding Author: zhaimingjie@ihep.ac.cn

TeV / 184

**Search for direct production of stau pairs in  $\sqrt{s} = 13$  TeV  $pp$  collisions with the ATLAS detector**

Corresponding Author: guolei@ihep.ac.cn

TeV / 185

**Search for lepton-flavour violation in high-mass dilepton final states using ATLAS run2 data**

Corresponding Author: luxin.zhang@cern.ch

TeV / 186

**Search for dark photons in rare Z boson decays with the ATLAS detector**

Corresponding Author: nh196245@mail.ustc.edu.cn

TeV / 187

**Search for a new  $Z'$  gauge boson in the  $\mu\mu\nu$  final state with the ATLAS detector**

TeV / 188

**A search for heavy right-handed Majorana neutrinos with ATLAS**

Corresponding Author: tzhao@cern.ch

Instrumental / 189

**Progress of the RPC development towards the production for ATLAS Phase-II upgrade**

Corresponding Author: dongshuo.du@cern.ch

Instrumental / 190

## **CMS-GEM upgrade progress and GEM assembly at Peking University**

**Corresponding Author:** 1801110097@pku.edu.cn

**Instrumental / 192**

## **CMS-GEM ME0 electronics board design, prototyping and production plan**

**Corresponding Author:** 2201110119@pku.edu.cn

**Instrumental / 193**

## **CMS iRPC Backend and Trigger status**

**Corresponding Author:** zhaojz@ihep.ac.cn

**Instrumental / 194**

## **Data analysis for iRPC BE/TRG beam test**

**Corresponding Author:** diaowz@ihep.ac.cn

**Instrumental / 195**

## **Grid computing for LHC experiments in China**

**Corresponding Author:** jiangxw@ihep.ac.cn

**Instrumental / 196**

## **Dark SHINE Simulation software framework**

**Corresponding Author:** xuliangz@sjtu.edu.cn

**Instrumental / 197**

## **IHEP ROC Status Report**

**Corresponding Author:** chenye@ihep.ac.cn

**Instrumental / 198**

## **Run3 CMS EGamma performance (Remote)**

**Corresponding Author:** akapoor@cern.ch

**Instrumental / 199**

## **Optimal transport solutions for pileup mitigation at hadron colliders (Remote)**

**Corresponding Author:** fabio.iemmi@cern.ch

**Instrumental / 200**

## **Ultimate calibration and performance of the CMS Electromagnetic Calorimeter in LHC Run 2(Remote)**

**Corresponding Author:** wangjin3@ihep.ac.cn

**Instrumental / 201**

## **RPC operation in the ATLAS experiment(Remote)**

**Corresponding Author:** mohamed.zaazoua@cern.ch

**Instrumental / 202**

## **Tau trigger scale factor studies**

**Corresponding Authors:** xiaohu.sun@cern.ch, botao.guo@stu.pku.edu.cn

**Instrumental / 203**

## **CMS DQM-DC operations and performance during Run3 data-taking**

**Corresponding Author:** tahir.javaid@cern.ch

**Instrumental / 204**

## **Constituent-based W-boson tagging with the ATLAS detector**

**Corresponding Author:** wangsd@ihep.ac.cn

**Instrumental / 205**

## **Performance of heavy flavour jet identification in boosted topologies with full Run2 data**

**Corresponding Author:** congqiao.li@cern.ch

**TeV / 206**

## **Observation of $WW\gamma$ production**

**Corresponding Author:** zhe.guan@cern.ch

**TeV / 207**

## **VBS $W\gamma$ at CMS**

**Corresponding Author:** xiaohu.sun@cern.ch

**TeV / 208**

## **$ZZ$ cross section measurement at 13.6TeV with the ATLAS detector**

**Corresponding Author:** singingstar@mail.ustc.edu.cn

**TeV / 209**

## **Evidence of pair-production of longitudinally polarized vector bosons in $ZZ$**

**Corresponding Author:** dongshuo.du@cern.ch

**TeV / 210**

## **Observation of $WZ\gamma$ production in pp collisions with ATLAS detector**

TeV / 211

## **Precise measurement of $Z\gamma$ +jets final state and search for neutral triple gauge couplings**

Corresponding Author: danningliu@sjtu.edu.cn

TeV / 212

## **Observation and measurement of the cross-sections of the electroweak and total production of a $Z\gamma$ pair in association with two jets in pp collisions at $\sqrt{s} = 13\text{TeV}$ with the ATLAS detector (remote)**

Corresponding Author: qibin.liu@sjtu.edu.cn

TeV / 213

## **Observation of Four-Top-Quark Production in the Multi-lepton Final State with the ATLAS Experiment (remote)**

Corresponding Author: brianchenfff@sjtu.edu.cn

TeV / 214

## **Search for heavy Higgs bosons decaying to a top quark pair with the ATLAS detector**

Corresponding Author: yizhou.cai@cern.ch

TeV / 215

## **Search for a new high-mass resonance decaying into $Z\gamma$ final state in pp collisions at $\sqrt{s} = 13\text{TeV}$ with the ATLAS detector**

Corresponding Author: fabio.lucio@cern.ch

TeV / 216

## **Search for leptoquarks decaying into the $b\tau$ final state in pp collisions at $\sqrt{s} = 13\text{TeV}$ with the ATLAS detector**

Corresponding Author: yimin.che@cern.ch

TeV / 217

**Search for the production of an excited bottom quark decaying to  $tW$  (leptonic top + hadronic  $W$ ) in pp collisions at 13 TeV (CMS)**

Corresponding Author: houbr@ihep.ac.cn

TeV / 218

**Search for a new heavy boson  $W'$  decaying to a top quark and a bottom quark**

Corresponding Author: yuanrui@sjtu.edu.cn

TeV / 219

**Search for  $W/Z \rightarrow W/Z$  production in a final state 2 with one lepton or two opposite-sign leptons using the full Run 2  $W/Z$  collisions data at  $\sqrt{s} = 13$  TeV**

Corresponding Author: mln517@sjtu.edu.cn

HF/HI/QCD / 221

**Observation and differential cross-section measurements of electroweak  $W/Z$  production in  $W/Z$  collisions at  $\sqrt{s} = 13$  TeV with the ATLAS detector (Remote)**

Corresponding Author: wangz1996@sjtu.edu.cn

HF/HI/QCD / 222

**Measurements of energy correlators and alphas extraction**

Corresponding Author: zhen.lin@cern.ch

HF/HI/QCD / 223

**First Direct Observation of Collider Neutrinos with FASER**

Corresponding Author: jinfeng.liu@cern.ch



HF/HI/QCD / 224

## **Study of isolated photon production with ALICE**

Corresponding Author: ran.xu@cern.ch

TeV / 225

## **First Results from the Search for Dark Photons with the FASER Detector at the L**

Corresponding Author: pangh19@mails.tsinghua.edu.cn

HF/HI/QCD / 226

## **Combination and summary of ATLAS dark matter searches interpreted in the context of 2HDM+a**

Corresponding Author: nkvu@sjtu.edu.cn

Poster Session / 227

## **b-tagging trigger calibration with ttbar event**

Corresponding Author: yuhui.miao@cern.ch

228

## **Improved Asymptotic Formulae for Statistical Interpretation Based on Likelihood**

Corresponding Author: ligang.xia@nju.edu.cn

Instrumental / 229

## **Status of Progress of CEPC Calorimeter R&D**

Corresponding Author: liuyong@ihep.ac.cn

Instrumental / 230

## **CEPC Flavor Physics**

**Corresponding Author:** calibbi@nankai.edu.cn

**Instrumental / 231**

## **Silicon detector at the CEPC**

**Corresponding Author:** liangzj@ihep.ac.cn

**Instrumental / 232**

## **CEPC New Physics Studies**

**Corresponding Author:** jialiu@pku.edu.cn

**Instrumental / 233**

## **Electroweak Physics at the CEPC**

**Corresponding Author:** jiyin\_gu@fudan.edu.cn

**Instrumental / 234**

## **Consideration on CEPC Electronics**

**Corresponding Author:** weiw@ihep.ac.cn

**Public Lecture / 235**

## **“小爆炸”：相对论重离子碰撞**

**Corresponding Author:** mayugang@fudan.edu.cn

**Poster Session / 236**

## **Performance of ATLAS Photon ID**

**Poster Session / 237**

## **Search for Higgs pairs in the bby final state with the ATLAS Experiment**

Poster Session / 238

## **Performance and Calibration of quark/gluon jet tagger and search for new resonances**

Corresponding Author: moony2628@sjtu.edu.cn

Poster Session / 239

## **Test results of Quality Control Test Structure of the USTC-IME LGAD pre-production**

Poster Session / 240

## **Improvement of jet reconstruction in forward region with time information from HGTD.**

Poster Session / 241

## **ACP technique for flip chip bonding**

Corresponding Author: aonan.wang@cern.ch

Poster Session / 242

## **LGAD based Beam monitor development**

Corresponding Author: fanyy@ihep.ac.cn

Poster Session / 243

## **Dark Photon search initiative at SHINE facility (Dark SHINE)**

Corresponding Author: ivychen8@sjtu.edu.cn

Poster Session / 244

## **Dark SHINE Simulation software framework**

**Corresponding Author:** xuliangz@sjtu.edu.cn

**Poster Session / 245**

## **R&D of Dark SHINE LYSO Crystal ECAL**

**Corresponding Author:** zhiyuzhao@sjtu.edu.cn

**Poster Session / 246**

## **R&D of Dark SHINE Strip Tracker**

**Corresponding Author:** lucker@sjtu.edu.cn

**Poster Session / 247**

## **R&D of Dark SHINE Hadronic Calorimeter (HCAL)**

**Corresponding Authors:** yuanrui@sjtu.edu.cn, jssduzcx@163.com, wangz1996@sjtu.edu.cn

**Poster Session / 248**

## **Development of readout electronics for Dark SHINE ECAL**

**Corresponding Authors:** wuweihaio@sjtu.edu.cn, tangjn@sjtu.edu.cn

**Poster Session / 249**

## **Simulation study of the muon flux produced by 1.6 GeV electrons in a beam**

**Corresponding Author:** liufangchao@sjtu.edu.cn

**Poster Session / 250**

## **Particle physics opportunities at the Shanghai SHINE facility with secondary**

**Corresponding Author:** kimsiang84@sjtu.edu.cn

**Poster Session / 251**

## **Development of a muon spin polarization monitor for the J-PARC Muon g-2/EDM experiment**

**Poster Session / 252**

## **Muon Lifetime Measurement with Run-1 data of Muon g-2 Experiment at Fermi**

**Corresponding Author:** lzj777@sjtu.edu.cn

**Poster Session / 253**

## **Development of a muon entrance trigger system for the PSI muEDM experiment**

**Corresponding Author:** hutianqi@sjtu.edu.cn

**Poster Session / 254**

## **Performance of a prototype PSI muEDM muon entrance detector measured with**

**Corresponding Author:** wong.gm@sjtu.edu.cn

**Poster Session / 255**

## **Reconstruction of Neutrino Events in TRIDENT Project based on GNN**

**Corresponding Author:** mo\_cen@sjtu.edu.cn