

Top Decay at Next-to-Next-to-Next-to-Leading Order in QCD

陈龙 (Long Chen)

山东大学物理学院 (Shandong University)

第九届中国 LHC 物理年会 (CLHCP2023)

2023 年 11 月 16-20 日, 上海

Based on: [2309.01937 and work in progress]
In collaboration with Xiang Chen, Xin Guan and Yan-Qing Ma



Why Study the Top Quark

- Heaviest fundamental particle in SM

$$m_t = 172.69 \pm 0.30 \text{ GeV}$$

- Precision test of SM mechanism, and pros for possible BSM physics
- Decay exclusively to $b + W$ before hadronization:

$$\Gamma_t = 1.4 \text{ GeV} \gg \Lambda_{\text{QCD}}$$


Standard Model of Elementary Particles

three generations of matter (fermions)			interactions / force carriers (bosons)	
mass $\sim 2.2 \text{ MeV/c}^2$	I	II	III	
charge $\frac{2}{3}$	u	c	t	g
spin $\frac{1}{2}$	up	charm	top	Higgs
mass $\sim 4.7 \text{ MeV/c}^2$				
charge $\frac{-1}{3}$	d	s	b	
spin $\frac{1}{2}$	down	strange	bottom	
mass $\sim 0.511 \text{ MeV/c}^2$				
charge -1	e	μ	τ	
spin $\frac{1}{2}$	electron	muon	tau	
mass $\sim 1.0 \text{ eV/c}^2$				
charge 0	ν_e	ν_μ	ν_τ	
spin $\frac{1}{2}$	electron neutrino	muon neutrino	tau neutrino	
mass $\sim 0.17 \text{ MeV/c}^2$				
charge 0				
spin $\frac{1}{2}$				
mass $\sim 173.1 \text{ GeV/c}^2$				
charge ± 1				
spin $\frac{1}{2}$				

SCALAR BOSONS

GAUGE BOSONS VECTOR BOSONS
Z boson
W boson
γ

- Convergence of the perturbative QCD series (e.g. renormalon issue)




Top Quark Mass m_t and Decay Width Γ_t

- PDG average for m_t :
 $172.69 \pm 0.30 \text{ GeV}$

- Current best measurement for Γ_t :


$$1.36 \pm 0.02 (\text{stat.})^{+0.14}_{-0.11} (\text{syst.}) \text{ GeV}$$

- Experimental uncertainties anticipated at future colliders:
 $20 \sim 26 \text{ MeV}$



The W-helicities in Top Decay

W from $t \rightarrow b + W^+ + X_{\text{QCD}}$ is polarized even if the t -quark is unpolarized



- The current best measurements: $f_0 = 0.684 \pm 0.005 \text{ (stat.)} \pm 0.014 \text{ (syst.)}$,
 $f_L = 0.318 \pm 0.003 \text{ (stat.)} \pm 0.008 \text{ (syst.)}$ and $f_R = -0.002 \pm 0.002 \text{ (stat.)} \pm 0.014 \text{ (syst.)}$.
- Notoriously difficult to be predicted theoretically to high precision

Much Theoretical Work Done So Far

Given the key role played by the top-quark both in SM precision test and searching for BSM, there have been vast amount of works done in literature regarding $t \rightarrow b + W^+ + X_{\text{QCD}}$.

● The inclusive Γ_t

- ▶ Up to NNLO in QCD: [Jezabek etc 88; Czarnecki etc 90; Li etc 90; Czarnecki etc 98; Chetyrkin etc 99; Fischer etc 01; Blokland etc 04'05;.....; Czarnecki etc 10; Meng etc 22; Chen etc 22]
- ▶ @NNNLO in QCD: [LC, Chen, Guan, Ma 23; Chen, Li, Li, Wang, Wang, Wu 23] [→ See also talk by Y.F. Wang]
[Datta, Rana, Ravindran, Sarkar 23 (only virtuals)]
- ▶ NLO Electroweak: [Denner Sack 91; Eilam, Mendel Migneron Soni 91]

● W-helicities $f_{L,R,0}$

- ▶ @NNLO in QCD: [Czarnecki, Korner, Piclum 10; Gao, Li, Zhu 12; Brucherseifer, Caola, Melnikov 13; Czarnecki, Groote Korner ,Piclum 18]
- ▶ @NNNLO in QCD: [LC, Chen, Guan, Ma 23]
- ▶ NLO Electroweak: [Do, Groote, Korner, Mauser 02]


● Differential results

- ▶ QCD:
 - @NLO [Fischer, Groote, Korner, Mauser 01; Brandenburg, Si, Uwer 02; Bernreuther, Gonzalez , Mellei 14; Kniehl, Nejad 21]
 - @NNLO [Gao, Li, Zhu 12; Brucherseifer, Caola, Melnikov 13; Campbell, Neumann, Sullivan 20]
 - @NNNLO in QCD: [LC, Chen, Guan, Ma 23]

Cut Diagrams for Top Decay Width

Γ_t in terms of the **semi-inclusive** $\mathcal{W}_{tb}^{\mu\nu}$

$$\Gamma_t = \frac{1}{2m_t} \int \frac{d^{d-1}\mathbf{k}}{(2\pi)^{d-1} 2E} \mathcal{W}_{tb}^{\mu\nu} \sum_{\lambda}^{L,R,0} \varepsilon_{\mu}^{*}(\mathbf{k}, \lambda) \varepsilon_{\nu}(\mathbf{k}, \lambda),$$



$$\begin{aligned} \mathcal{W}_{tb}^{\mu\nu}(p, \mathbf{k}) &= W_1 g^{\mu\nu} + W_2 p^{\mu} p^{\nu} + W_3 \mathbf{k}^{\mu} \mathbf{k}^{\nu} \\ &+ W_4 (p^{\mu} \mathbf{k}^{\nu} + \mathbf{k}^{\mu} p^{\nu}) + W_5 i \epsilon^{\mu\nu\rho\sigma} p_{\rho} \mathbf{k}_{\sigma}, \end{aligned}$$

Selection Criteria: the **cut diagrams** of t -quark self-energy function with exactly one **(cut) W propagator** interacting with the *external* t -quark plus (up to 3) QCD loops

Loop and Phase-space Integration

- Loop integrals are reduced using IBP [Chetyrkin 81] relations done with Blade [Guan, Liu, Ma 20], and the resulting *masters* are calculated using DE method [Kotikov 90; Remiddi 97] with AMFlow [Liu, Ma 22]
- The phase-space integrals, except for **W-momentum k** , are treated in the same manner as loop integrals by means of the reverse unitarity [Anastasiou, Melnikov 02]
- The **IR-divergent** phase-space integration of $\mathcal{W}_{tb}^{\mu\nu}$ over k are done “*manually*” using its power-log series representation (PSE) with ϵ assigned with **non-zero** numbers.

► Level of Complexity:

7×10^4 integrals reduced to 2988 master integrals, for which PSE about 200 orders in k_0 are derived with the above method.

► Consistency Check:

A perfect agreement in the result for the **inclusive** Γ_t with $\frac{d^{d-1}k}{(2\pi)^{d-1}2E}$ calculated in this way and by directly applying the reverse unitarity [Anastasiou, Melnikov 02]

Results for the Inclusive Γ_t

The QCD effects on Γ_t in SM can be parameterized as

$$\Gamma_t = \Gamma_0 \left[\mathbf{c}_0 + \frac{\alpha_s}{\pi} \mathbf{c}_1 + \left(\frac{\alpha_s}{\pi} \right)^2 \mathbf{c}_2 + \left(\frac{\alpha_s}{\pi} \right)^3 \mathbf{c}_3 + \mathcal{O}(\alpha_s^4) \right],$$

with $\Gamma_0 \equiv \frac{G_F m_W^2 m_t |V_{tb}|^2}{12\sqrt{2}}$.

We choose $\mu = m_t/2$, motivated by the **kinetic energy** $m_t - m_W - m_b$ of the QCD radiations, at which our N3LO result reads: [LC, Chen, Guan, Ma 23]

$$\begin{aligned}\Gamma_t &= 1.48642 \textcolor{blue}{-} 0.140877 \textcolor{green}{-} 0.023306 \textcolor{red}{-} 0.007240 \text{ GeV} \\ &= \mathbf{1.31500} \text{ GeV}\end{aligned}$$


SM Inputs:

$$\begin{aligned}G_F &= 1.166379 \times 10^{-5} \text{ GeV}^{-2}, & m_t &= 172.69 \text{ GeV}, \\ m_W &= 80.377 \text{ GeV}, & \alpha_s(m_t/2) &\approx 0.1189.\end{aligned}$$

The **leading-color** part of Γ_t agrees with a parallel computation [Chen, Li, Li, Wang, Wang, Wu 23]

The QCD Scale Uncertainty of Γ_t

The scale dependence of the fixed-order results for Γ_t in $\mu/m_t \in [0.1, 1]$



- NNLO scale-variation **never** cover the NNNLO result at any scales less than $\mu/m_t = 0.6$.
- Pure $\mathcal{O}(\alpha_s^3)$ correction decreases Γ_t by $\sim 0.8\%$ of the NNLO result at $\mu = m_t$ roughly 10 MeV(exceeding NNLO scale-hand)

The Offshell W and Finite m_b Effects

- With $\frac{1}{k^2 - m_W^2 + i\epsilon} \rightarrow \frac{1}{k^2 - m_W^2 + im_W \Gamma_W}$,

$$\begin{aligned}\Gamma_t(m_W) \rightarrow \tilde{\Gamma}_t &= \int_0^{m_t^2} \frac{dk^2}{2\pi} \frac{2m_W \Gamma_W}{(k^2 - m_W^2)^2 + (m_W \Gamma_W)^2} \Gamma_t(m_W^2 \rightarrow k^2) \\ &= \tilde{\Gamma}_0 \left[\tilde{\mathbf{c}}_0 + \frac{\alpha_s}{\pi} \mathbf{c}_1 + \left(\frac{\alpha_s}{\pi}\right)^2 \mathbf{c}_2 + \left(\frac{\alpha_s}{\pi}\right)^3 \mathbf{c}_3 + \mathcal{O}(\alpha_s^4) \right],\end{aligned}$$

we find: $\frac{\tilde{\mathbf{c}}_i - \mathbf{c}_i}{\mathbf{c}_i}$ takes $-1.54\%, -1.53\%, -1.39\%, -1.23\%$ for $i = 0, 1, 2, 3$.

- Similarly, keeping $m_b = 4.78 \text{ GeV}$, we find: $\frac{\mathbf{c}_1^{m_b} - \mathbf{c}_1}{\mathbf{c}_1} \approx \frac{\mathbf{c}_2^{m_b} - \mathbf{c}_2}{\mathbf{c}_2} \approx -1.47\%$.
- The NLO electroweak K-factor is re-evaluated to be $K_{EW}^{NLO} = 1.0168$.

Taking these misc-effects into account, we finally obtain the **to-date most-precise high-precision theoretical prediction**:

$$\boxed{\Gamma_t = 1.3148^{+0.003}_{-0.005} \times |V_{tb}|^2 + 0.027 (m_t - 172.69) \text{ GeV}}$$

the error of which meets the request by future colliders.

Results for W-helicity Fractions

Top decay width with polarized W :

$$\Gamma_{\lambda} = \frac{1}{2m_t} \int \frac{d^{d-1}k}{(2\pi)^{d-1} 2E} \mathcal{W}_{tb}^{\mu\nu} \varepsilon_{\mu}^{*}(k, \lambda) \varepsilon_{\nu}(k, \lambda)$$

The **W -helicity fractions** $f_{\lambda}^{[n]} = \frac{\sum_{i=0}^n \Gamma_{\lambda}^{[n]}}{\sum_{i=0}^n \Gamma_t^{[n]}}$ truncated to $\mathcal{O}(\alpha_s^3)$ in massless QCD:

$$f_0^{[3]} = 0.697706 - 0.008401 - 0.001954 - 0.000613,$$

$$= 0.686737,$$

$$f_L^{[3]} = 0.302294 + 0.007254 + 0.001799 + 0.000586,$$

$$= 0.311933,$$

$$f_R^{[3]} = 0. + 0.001147 + 0.000155 + 0.000027,$$

$$= 0.001330,.$$


(The above results evaluated at $\mu = m_t$ agree with [Czarnecki, Korner, Piclum 10] up to NNLO)

With NLO EW-correction and m_b effects included, our **final results** read:

$$f_0^{[3]} = 0.686_{-0.003}^{+0.002}, \quad f_L^{[3]} = 0.312_{-0.002}^{+0.001}, \quad f_R^{[3]} = 0.00157_{-0.00002}^{+0.00002}.$$


Results for $\cos \theta^*$ Angular Distribution

$$\frac{1}{\Gamma_t} \frac{d\Gamma_t}{d \cos \theta^*} = \frac{3}{4} (\sin^2 \theta^*) f_0 + \frac{3}{8} (1 - \cos \theta^*)^2 f_L + \frac{3}{8} (1 + \cos \theta^*)^2 f_R$$



where θ^* is the angle between the charged-lepton momentum from the W-decay in W-rest frame and the W-momentum in t-rest frame.


Results for W-energy Distribution



- ▶ **In the bulk:** QCD corrections are **positive and quite sizable**: pure $\mathcal{O}(\alpha_s^3)$ correction modifies the lowest order by $7 \sim 14\%$ for $E \in [94, 104]$ GeV.
- ▶ **In the rightmost 1 GeV-bin:** QCD corrections up to $\mathcal{O}(\alpha_s^3)$ decrease the Born-level result.

Bonus: lepton-pair invariant-mass spectrum in B-decay

$$\Gamma(b \rightarrow ul\bar{v}_l) = \Gamma_0 \left(1 - 2.4131 \left(\frac{\alpha_s}{\pi} \right) - 21.27 \left(\frac{\alpha_s}{\pi} \right)^2 - 270.7 \left(\frac{\alpha_s}{\pi} \right)^3 \right).$$



- ▶ The leading-color inclusive part agrees with a parallel computation [Chen, Li, Li, Wang, Wang, Wu 23].
- ▶ The $\mathcal{O}(\alpha_s^3)$ corrections agrees well with a recent approximation [Fael, Usovitsch 23].

Summary and Outlook

- ✓ We have provided the **to-date most-precise high-precision theoretical prediction** for top-quark decay width:

$$\Gamma_t = 1.3148^{+0.003}_{-0.005} \times |V_{tb}|^2 + 0.027 (m_t - 172.69) \text{ GeV}$$

the error of which meets the request by future colliders.

- ✓ By a novel approach to complete IR-divergent phase-space integration over W -momentum, we determined, in addition, W -helicity fractions, $\cos \theta^*$ distribution and W -energy distribution at α_s^3 for the first time.
- ✓ Furthermore, the lepton invariant-mass distribution in the B-meson decay $b \rightarrow u l \bar{\nu}_l$ is derived up to α_s^3 .
- ✓ The approach can be readily applied to the decay of polarized t -quarks at α_s^3 , as well as the mixed QCD-electroweak corrections.

Summary and Outlook

- ✓ We have provided the **to-date most-precise high-precision theoretical prediction** for top-quark decay width:

$$\Gamma_t = 1.3148^{+0.003}_{-0.005} \times |V_{tb}|^2 + 0.027 (m_t - 172.69) \text{ GeV}$$

the error of which meets the request by future colliders.

- ✓ By a novel approach to complete IR-divergent phase-space integration over W -momentum, we determined, in addition, W -helicity fractions, $\cos \theta^*$ distribution and W -energy distribution at α_s^3 for the first time.
- ✓ Furthermore, the lepton invariant-mass distribution in the B -meson decay $b \rightarrow u l \bar{v}_l$ is derived up to α_s^3 .
- ✗ The approach can be readily applied to the decay of polarized t -quarks at α_s^3 , as well as the mixed QCD-electroweak corrections.

谢谢 (Thank you)!