

第九届中国LHC物理年会

The 9th China LHC Physics Workshop

Recent charm baryon spectroscopy results at LHCb

Zhenhong Wu, Zhihao Xu




中国科学院大学

University of Chinese Academy of Sciences







CLHCP 2023
Nov. 16 to 20, 2023



Motivation - I




- Why we do spectroscopy?
- Missing particles in hadron multiplets;
 - Ground QQq , QQQ states
 - Excited Qqq states
- Complex inner structure of excited states;
 - Molecular? Pentaquark? (hidden $q\bar{q}$?)
 - Radially mode? Orbital mode?



Motivation - II

- What can we do?
- Searching for new decays.
- Expanding known decays.
- Mass, width (lifetime);
- Production;
- Branching ratio;
- Quantum numbers ($I^G J^{PC}$).



patrick.koppenburg@cern.ch 2023-02-10

LHCb experiment - I

$$\sigma_{\text{IP}} = 20 \text{ } \mu\text{m}$$


$$\sigma_{\tau} = 45 \text{ fs}$$

$$\sigma_p/p \sim 0.5\% - 1.0\%$$

$$\sigma_E/E = \frac{10\%}{\sqrt{E}} \pm 1\%$$

$$\epsilon(K \rightarrow K) \sim 95\%$$


$$\text{Mis-ID } \epsilon(\pi \rightarrow K) \sim 5\%$$



- LHCb designed for study charmed and beauty hadron.
- Excellent vertex, tracking and PID performance.

LHCb experiment - II

- LHCb collected the largest samples of reconstructed heavy hadrons during LHC Run 1 and Run 2.




patrick.koppenburg@cern.ch 2023-02-10

[LHCb collaboration, P. Koppenburg, List of hadrons observed at the LHC, [LHCb-FIGURE-2021-001](#), 2021, and [2023 updates](#)]


Zhihao Xu(UCAS) @CLHCP2023

Excited Ω_c^0 states - I

- Starting of the story...
- In 2017, LHCb observed five new excited Ω_c^0 in $m(\Xi_c^+ K^-)$.
- **Four** of them confirmed by Belle.



2017 LHCb result
[PHYS. REV. LETT. 118 \(2017\) 182001](#)




2018 Belle result
[PHYS. REV. D 97 \(2018\) 5, 051102](#)

Excited Ω_c^0 states - II

- In 2021, LHCb observed **four** of them from $\Omega_b^- \rightarrow \Omega_c^{(**)0} \pi^-$
- Spin test done, but no solid conclusion.

[PHYS. REV. D 104 \(2021\) 9, 9.](#)



State	Observable	Measurement
Ω_b^-	m	$6044.3 \pm 1.2 \pm 1.1^{+0.19}_{-0.22}$ MeV
	$\mathcal{R}$	$1.35 \pm 0.11 \pm 0.05$
Threshold structure	Significance	Stat. 4.3 σ Sys.
$\Omega_c(3000)^0$	Significance	6.2 σ
	ΔM	$37.6 \pm 0.9 \pm 0.9$ MeV
	m	$2999.2 \pm 0.9 \pm 0.9^{+0.19}_{-0.22}$ MeV
	Γ	$4.8 \pm 2.1 \pm 2.5$ MeV
	$\mathcal{P}$	$0.11 \pm 0.02 \pm 0.04$
$\Omega_c(3050)^0$	J rejection	0.5 σ ($J = 1/2$), 0.8 σ ($J = 3/2$), 0.4 σ ($J = 5/2$)
	Significance	9.9 σ
	ΔM	$88.5 \pm 0.3 \pm 0.2$ MeV
	m	$3050.1 \pm 0.3 \pm 0.2^{+0.19}_{-0.22}$ MeV
	Γ	< 1.6 MeV, 95% CL
$\Omega_c(3065)^0$	$\mathcal{P}$	$0.15 \pm 0.02 \pm 0.02$
	J rejection	2.2 σ ($J = 1/2$), 0.1 σ ($J = 3/2$), 1.2 σ ($J = 5/2$)
	Significance	11.9 σ
	ΔM	$104.3 \pm 0.4 \pm 0.4$ MeV
	m	$3065.9 \pm 0.4 \pm 0.4^{+0.19}_{-0.22}$ MeV
$\Omega_c(3090)^0$	Γ	$1.7 \pm 1.0 \pm 0.5$ MeV
	$\mathcal{P}$	$0.23 \pm 0.02 \pm 0.02$
	J rejection	3.6 σ ($J = 1/2$), 0.6 σ ($J = 3/2$), 1.2 σ ($J = 5/2$)
	Significance	7.8 σ
	ΔM	$129.4 \pm 1.1 \pm 1.0$ MeV
$\Omega_c(3120)^0$	m	$3091.0 \pm 1.1 \pm 1.0^{+0.19}_{-0.22}$ MeV
	Γ	$7.4 \pm 3.1 \pm 2.8$ MeV
	$\mathcal{P}$	$0.19 \pm 0.02 \pm 0.04$
	J rejection	0.3 σ ($J = 1/2$), 0.8 σ ($J = 3/2$), 0.5 σ ($J = 5/2$)
	$\mathcal{P}$	< 0.03, 95% CL


New excited Ω_c^0 states - I

PHYS. REV. LETT. 131 (2023) 131902.

- **New states observed:** $\Omega_c(3185)^0$ and $\Omega_c(3327)^0$
- **All previous states confirmed**, and masses and widths measured with the highest precision.
- Detailed study of the threshold enhancement.
- Detailed study of the $\Omega_c(3185)^0$ with alternative model. } as systematics

Resonance	m (MeV)	Γ (MeV)
$\Omega_c(3000)^0$	$3000.44 \pm 0.07^{+0.07}_{-0.13} \pm 0.23$	$3.83 \pm 0.23^{+1.59}_{-0.29}$
$\Omega_c(3050)^0$	$3050.18 \pm 0.04^{+0.06}_{-0.07} \pm 0.23$	$0.67 \pm 0.17^{+0.64}_{-0.72}$
		< 1.8 MeV, 95% C.L.
$\Omega_c(3065)^0$	$3065.63 \pm 0.06^{+0.06}_{-0.06} \pm 0.23$	$3.79 \pm 0.20^{+0.38}_{-0.47}$
$\Omega_c(3090)^0$	$3090.16 \pm 0.11^{+0.06}_{-0.10} \pm 0.23$	$8.48 \pm 0.44^{+0.61}_{-1.62}$
$\Omega_c(3119)^0$	$3118.98 \pm 0.12^{+0.09}_{-0.23} \pm 0.23$	$0.60 \pm 0.63^{+0.90}_{-1.05}$
		< 2.5 MeV, 95% C.L.
$\Omega_c(3185)^0$	$3185.1 \pm 1.7^{+7.4}_{-0.9} \pm 0.2$	$50 \pm 7^{+10}_{-20}$
$\Omega_c(3327)^0$	$3327.1 \pm 1.2^{+0.1}_{-1.3} \pm 0.2$	$20 \pm 5^{+13}_{-1}$


$\Omega_c(3000)^0 \rightarrow \Xi_c^+ K^-$	$\Omega_c(3065)^0 \rightarrow \Xi_c^+ (\rightarrow \Xi_c^+ \gamma) K^-$
$\Omega_c(3050)^0 \rightarrow \Xi_c^+ K^-$	$\Omega_c(3090)^0 \rightarrow \Xi_c^+ (\rightarrow \Xi_c^+ \gamma) K^-$
$\Omega_c(3065)^0 \rightarrow \Xi_c^+ K^-$	$\Omega_c(3119)^0 \rightarrow \Xi_c^+ (\rightarrow \Xi_c^+ \gamma) K^-$
$\Omega_c(3090)^0 \rightarrow \Xi_c^+ K^-$	$\Omega_c(3185)^0 \rightarrow \Xi_c^+ K^-$
$\Omega_c(3119)^0 \rightarrow \Xi_c^+ K^-$	$\Omega_c(3327)^0 \rightarrow \Xi_c^+ K^-$




New

New excited Ω_c^0 states - II


- Three $m(\Xi_c^+ K^-)$ studies from LHCb...
- And one $m(\Xi_c^+ K^-)$ study from Belle...
- What happens on $\Omega_c(3119)^0$?




PHYS. REV. LETT. 118 (2017) 182001



PHYS. REV. LETT. 131 (2023) 131902.



PHYS. REV. D 104 (2021) 9, 9.




New excited Ω_c^0 states - III



Excited Ξ_c^0 states - I

[PHYS. REV. LETT. 124 \(2020\) 222001](#)

- Three excited Ξ_c^0 were observed in prompt $m(\Lambda_c^+ K^-)$.
- Using LHCb 2016-2018 data, at 13 TeV and 5.4 fb^{-1} .



Resonance	Peak of ΔM [MeV]	Mass [MeV]			Γ [MeV]
$\Xi_c(2923)^0$	$142.91 \pm 0.25 \pm 0.20$	$2923.04 \pm 0.25 \pm 0.20 \pm 0.14$			$7.1 \pm 0.8 \pm 1.8$
$\Xi_c(2939)^0$	$158.45 \pm 0.21 \pm 0.17$	$2938.55 \pm 0.21 \pm 0.17 \pm 0.14$			$10.2 \pm 0.8 \pm 1.1$
$\Xi_c(2965)^0$	$184.75 \pm 0.26 \pm 0.14$	$2964.88 \pm 0.26 \pm 0.14 \pm 0.14$			$14.1 \pm 0.9 \pm 1.3$
		Stat.	Sys.	input	


Excited Ξ_c^0 states - II

PHYSICAL REVIEW D 108, 012020 (2023)

- $B^- \rightarrow \Lambda_c^+ \Lambda_c^- K^-$ decay is studied.
- Confirmed $\Xi_c(2923)^0$ and $\Xi_c(2939)^0$, consistent with prompt result.
- **Evidence of $\Xi_c(2880)^0 \rightarrow \Lambda_c^+ K^-$ observed (3.8σ).**
- No structure on $m(\Lambda_c^+ \Lambda_c^-)$ and $m(\Lambda_c^- K^-)$

State	Mass (MeV)	Width (MeV)
$\Xi_c(2880)^0$	$2881.8 \pm 3.1 \pm 8.5$	$12.4 \pm 5.2 \pm 5.8$
$\Xi_c(2923)^0$	$2924.5 \pm 0.4 \pm 1.1$	$4.8 \pm 0.9 \pm 1.5$
$\Xi_c(2939)^0$	$2938.5 \pm 0.9 \pm 2.3$	$11.0 \pm 1.9 \pm 7.5$

$$\frac{\mathcal{B}(B^- \rightarrow \Lambda_c^+ \Lambda_c^- K^-)}{\mathcal{B}(B^- \rightarrow D^+ D^- K^-)} = 2.36 \pm 0.11 \pm 0.22 \pm 0.25$$





Excited Ξ_b^- states

PHYS. REV. LETT. 126, 252003

- $\Xi_b(6100)^-$ observed by CMS in $m(\Xi_b^- \pi^+ \pi^-)$
- Two channels used to reconstruct Ξ_b^-

$$m(\Xi_b(6100)^-) = 6100 \pm 0.2(\text{stat.}) \pm 0.1(\text{syst}) \pm 0.6 (\Xi_b^-) \text{ MeV}$$

$$\Gamma(\Xi_b(6100)^-) < 1.9 \text{ MeV, C.L. 95\%}$$






New

New excited Ξ_b^0 states


PHYS. REV. LETT. 131 (2023) 171901

- Observation of two new excited Ξ_b^0 baryons: $\Xi_b(6087)^0, \Xi_b(6095)^0$.
- One state is confirmed: $\Xi_b(6100)^-$.
- First observation of $\Xi_b^0 \rightarrow \Xi_c^+ \pi \pi \pi$
- Best measurement on known $\Xi_b'^-$ and Ξ_b^{*-} states.

Q_0 ($\Xi_b^0(6087)$)	$16.20 \pm 0.20 \pm 0.06$
Γ ($\Xi_b^0(6087)$)	$2.43 \pm 0.51 \pm 0.10$
m_0 ($\Xi_b^0(6087)$)	$6087.24 \pm 0.20 \pm 0.06 \pm 0.5$ (Ξ_b^0)
Q_0 ($\Xi_b^0(6095)$)	$24.32 \pm 0.15 \pm 0.03$
Γ ($\Xi_b^0(6095)$)	$0.50 \pm 0.33 \pm 0.11$
m_0 ($\Xi_b^0(6095)$)	$6095.36 \pm 0.15 \pm 0.03 \pm 0.5$ (Ξ_b^0)





$$\Xi_{cc}^{++} \rightarrow \Xi_c'^+ \pi^+$$




JHEP 05 (2022) 038

- $\Xi_{cc}^{++} \rightarrow \Xi_c'^+ \pi^+$ observed with LHCb run-2 data.
- $\Xi_{cc}^{++} \rightarrow \Xi_c'^+ \pi^+$ is reconstructed partially.

$$\frac{\mathcal{B}(\Xi_{cc}^{++} \rightarrow \Xi_c'^+ \pi^+)}{\mathcal{B}(\Xi_{cc}^{++} \rightarrow \Xi_c^+ \pi^+)} = 1.41 \pm 0.17 \pm 0.10$$




$$\Xi_{bc}^+ \rightarrow J/\psi \Xi_c^+$$




CHINESE PHYS. C 47 (2023) 093001



- First search for $\Xi_{bc}^+ \rightarrow J/\psi \Xi_c^+$, with 9 fb^{-1} .
- Two peaking structures seen
 - Mass: 6571 MeV. Local (global) significance: $4.3(2.8) \sigma$
 - Mass: 6694 MeV. Local (global) significance: $4.1(2.4) \sigma$



$$\Xi_{bc}^+ \rightarrow J/\psi \Xi_c^+$$



$$\mathcal{R} = \frac{\sigma(\Xi_{bc}^+) \times \mathcal{B}(\Xi_{bc}^+ \rightarrow J/\psi \Xi_c^+) \times \mathcal{B}(\Xi_c^+ \rightarrow p K^- \pi^+)}{\sigma(B_c^+) \times \mathcal{B}(B_c^+ \rightarrow J/\psi D_s^+) \times \mathcal{B}(D_s^+ \rightarrow K^+ K^- \pi^+)}$$



Summary



- Heavy baryons
 - Observation of excited Ξ_c^0 baryons
 - Observation of excited Ξ_b^- baryons
 - **Observation of excited Ω_c^0 baryons (recent)**
 - **Observation of excited Ξ_b^0 baryons (recent)**
- Double Heavy baryons
 - Measurement of $\frac{\mathcal{B}(\Xi_{cc}^{++} \rightarrow \Xi_c'^+ \pi^+)}{\mathcal{B}(\Xi_{cc}^{++} \rightarrow \Xi_c^+ \pi^+)}$
 - First search for $\Xi_{bc}^+ \rightarrow J/\psi \Xi_c^+$

Future

- Higher luminosity
- Better detector
- Improved techniques
-
- More particles...
- More decays...
- Higher precision...
- Fulfill hadron picture



THANKS!



backup