

CEPC EW physics: towards White Paper

Jiayin Gu (顾嘉荫)

Fudan University

CLHCP2023
Nov 18, 2023

mainly based on an earlier draft of the EW white paper
and the CEPC Snowmass report [2205.08553]



The current status of the CEPC EW white paper

- ▶ **An earlier draft (last updated in 2019) with outdated run scenarios**
 - ▶ Z-pole measurements
 - ▶ W mass measurements (threshold and kinematic reconstruction)
 - ▶ Oblique parameter fit
 - ▶ SMEFT fit
 - ▶ New physics implications (Natural SUSY)
- ▶ **The CEPC Snowmass report [2205.08553]**
 - ▶ Updated measurement inputs
 - ▶ Updated SMEFT fit
 - ▶ New physics implications (see the previous talk by Jia)


Change of run scenarios

- ▶ **Z pole:** $\sim 8 \text{ ab}^{-1}?$ $\rightarrow 100 \text{ ab}^{-1}$
 - ▶ Many measurements are dominated by systematics, but A_e and A_τ from final state tau polarization measurements are significantly improved. (They were already considered in the earlier draft but were not official...)
- ▶ **WW threshold:** 3.2 ab^{-1} $\rightarrow 6 \text{ ab}^{-1}$
 - ▶ W mass: 1 MeV $\rightarrow 0.5 \text{ MeV}$ (We also got more optimistic?)
- ▶ **240 GeV:** 5.6 ab^{-1} $\rightarrow 20 \text{ ab}^{-1}$
 - ▶ Higgs and diboson ($e^+e^- \rightarrow WW$) measurements
- ▶ **Top threshold run:** no \rightarrow yes
 - ▶ The top mass measurement is an important input for EW fits!



Observable	current precision	CEPC precision (Stat. Unc.)	CEPC runs	main systematic
Δm_Z	2.1 MeV [37–41]	0.1 MeV (0.005 MeV)	Z threshold	E_{beam}
$\Delta \Gamma_Z$	2.3 MeV [37–41]	0.025 MeV (0.005 MeV)	Z threshold	E_{beam}
Δm_W	9 MeV [42–46]	0.5 MeV (0.35 MeV)	WW threshold	E_{beam}
$\Delta \Gamma_W$	49 MeV [46–49]	2.0 MeV (1.8 MeV)	WW threshold	E_{beam}
Δm_t	0.76 GeV [50]	$\mathcal{O}(10)$ MeV ^a	$t\bar{t}$ threshold	
ΔA_e	4.9×10^{-3} [37, 51–55]	1.5×10^{-5} (1.5×10^{-5})	Z pole ($Z \rightarrow \tau\tau$)	Stat. Unc.
ΔA_μ	0.015 [37, 53]	3.5×10^{-5} (3.0×10^{-5})	Z pole ($Z \rightarrow \mu\mu$)	point-to-point Unc.
ΔA_τ	4.3×10^{-3} [37, 51–55]	7.0×10^{-5} (1.2×10^{-5})	Z pole ($Z \rightarrow \tau\tau$)	tau decay model
ΔA_b	0.02 [37, 56]	20×10^{-5} (3×10^{-5})	Z pole	QCD effects
ΔA_c	0.027 [37, 56]	30×10^{-5} (6×10^{-5})	Z pole	QCD effects
$\Delta \sigma_{had}$	37 pb [37–41]	2 pb (0.05 pb)	Z pole	luminosity
δR_b^0	0.003 [37, 57–61]	0.0002 (5×10^{-6})	Z pole	gluon splitting
δR_c^0	0.017 [37, 57, 62–65]	0.001 (2×10^{-5})	Z pole	gluon splitting
δR_e^0	0.0012 [37–41]	2×10^{-4} (3×10^{-6})	Z pole	E_{beam} and t channel
δR_μ^0	0.002 [37–41]	1×10^{-4} (3×10^{-6})	Z pole	E_{beam}
δR_τ^0	0.017 [37–41]	1×10^{-4} (3×10^{-6})	Z pole	E_{beam}
δN_ν	0.0025 [37, 66]	2×10^{-4} (3×10^{-5})	ZH run ($\nu\nu\gamma$)	Calo energy scale

- ▶ Δ : absolute uncertainties, δ : relative uncertainties
- ▶ The constraints on A_e and A_τ mainly come from $e^+e^- \rightarrow \tau^+\tau^-$ with final state tau polarization measurements.
- ▶ A_μ , A_b and A_c are derived from the A^{FB} measurements and A_e .
 - ▶ Best way to present the results?

S & T parameters (earlier draft)




S & T parameters (earlier draft)



- What's the impact of the m_t measurement?

The Standard Model Effective Field Theory



+

X^3		ψ^4 and $\psi^2 D^2$	$\psi^3 \psi$	(LL)(LL)	(RR)(RR)	(LR)(RR)	
Q_{G1}	$f^{ABC} G^A_{\mu\nu} G^B_{\nu\rho} G^C_{\rho\mu}$	Q_{ψ^4}	$(\bar{\psi}\psi)^4$	Q_{LL}	$(\bar{\psi}_L\psi_L)(\bar{\psi}_L\psi_L)$	Q_{LR}	$(\bar{\psi}_L\psi_L)(\bar{\psi}_R\psi_R)$
Q_{G2}	$f^{ABC} \tilde{G}^A_{\mu\nu} G^B_{\nu\rho} G^C_{\rho\mu}$	$Q_{\psi^2 D^2}$	$(\bar{\psi}\psi)(\Box\bar{\psi}\psi)$	$Q_{LL}^{(1)}$	$(\bar{\psi}_L\psi_L)(\bar{\psi}_L\psi_L)$	$Q_{LR}^{(1)}$	$(\bar{\psi}_L\psi_L)(\bar{\psi}_R\psi_R)$
Q_{G3}	$f^{ABC} W^A_{\mu\nu} W^B_{\nu\rho} W^C_{\rho\mu}$	$Q_{\psi^2 D^2}$	$(\bar{\psi}^c D_\mu\psi)^2$	$Q_{LL}^{(2)}$	$(\bar{\psi}_L\psi_L)(\bar{\psi}_L\psi_L)$	$Q_{LR}^{(2)}$	$(\bar{\psi}_L\psi_L)(\bar{\psi}_R\psi_R)$
Q_{G4}	$f^{ABC} \tilde{W}^A_{\mu\nu} W^B_{\nu\rho} W^C_{\rho\mu}$	$Q_{\psi^2 D^2}$	$(\bar{\psi}^c D_\mu\psi)$	$Q_{LL}^{(3)}$	$(\bar{\psi}_L\psi_L)(\bar{\psi}_L\psi_L)$	$Q_{LR}^{(3)}$	$(\bar{\psi}_L\psi_L)(\bar{\psi}_R\psi_R)$
	$X^2\psi^2$	$\psi^2 X\psi$	$\psi^2\psi D$	$Q_{LL}^{(4)}$	$(\bar{\psi}_L\psi_L)(\bar{\psi}_L\psi_L)$	$Q_{LR}^{(4)}$	$(\bar{\psi}_L\psi_L)(\bar{\psi}_R\psi_R)$
Q_{G5}	$\bar{\psi}\psi G^A_{\mu\nu} G^A_{\nu\rho} G^A_{\rho\mu}$	$Q_{\psi^2 X\psi}$	$(\bar{\psi}\psi\psi)_\mu \psi^\mu W^A_\nu W^{\nu A}$	$Q_{LL}^{(5)}$	$(\bar{\psi}_L\tilde{D}_\mu\psi)(\bar{\psi}_L\psi_L)$	$Q_{LR}^{(5)}$	$(\bar{\psi}_L\tilde{D}_\mu\psi)(\bar{\psi}_R\psi_R)$
Q_{G6}	$\bar{\psi}\psi G^A_{\mu\nu} G^A_{\nu\rho} G^A_{\rho\mu}$	$Q_{\psi^2 X\psi}$	$(\bar{\psi}\psi\psi)_\mu \psi^\mu B_\nu W^{\nu A}$	$Q_{LL}^{(6)}$	$(\bar{\psi}_L\tilde{D}_\mu\psi)(\bar{\psi}_L\psi_L)$	$Q_{LR}^{(6)}$	$(\bar{\psi}_L\tilde{D}_\mu\psi)(\bar{\psi}_R\psi_R)$
Q_{G7}	$\bar{\psi}\psi W^A_{\mu\nu} W^A_{\nu\rho} W^A_{\rho\mu}$	$Q_{\psi^2 X\psi}$	$(\bar{\psi}\psi\psi)_\mu \psi^\mu B_\nu W^{\nu A}$	$Q_{LL}^{(7)}$	$(\bar{\psi}_L\tilde{D}_\mu\psi)(\bar{\psi}_L\psi_L)$	$Q_{LR}^{(7)}$	$(\bar{\psi}_L\tilde{D}_\mu\psi)(\bar{\psi}_R\psi_R)$
Q_{G8}	$\bar{\psi}\psi W^A_{\mu\nu} W^A_{\nu\rho} W^A_{\rho\mu}$	$Q_{\psi^2 X\psi}$	$(\bar{\psi}\psi\psi)_\mu \psi^\mu B_\nu W^{\nu A}$	$Q_{LL}^{(8)}$	$(\bar{\psi}_L\tilde{D}_\mu\psi)(\bar{\psi}_L\psi_L)$	$Q_{LR}^{(8)}$	$(\bar{\psi}_L\tilde{D}_\mu\psi)(\bar{\psi}_R\psi_R)$
Q_{G9}	$\bar{\psi}\psi B_\mu B^\mu$	$Q_{\psi^2 X\psi}$	$(\bar{\psi}\psi\psi)_\mu \psi^\mu B_\nu W^{\nu A}$	$Q_{LL}^{(9)}$	$(\bar{\psi}_L\tilde{D}_\mu\psi)(\bar{\psi}_L\psi_L)$	$Q_{LR}^{(9)}$	$(\bar{\psi}_L\tilde{D}_\mu\psi)(\bar{\psi}_R\psi_R)$
Q_{G10}	$\bar{\psi}\psi B_\mu B^\mu$	$Q_{\psi^2 X\psi}$	$(\bar{\psi}\psi\psi)_\mu \psi^\mu B_\nu W^{\nu A}$	$Q_{LL}^{(10)}$	$(\bar{\psi}_L\tilde{D}_\mu\psi)(\bar{\psi}_L\psi_L)$	$Q_{LR}^{(10)}$	$(\bar{\psi}_L\tilde{D}_\mu\psi)(\bar{\psi}_R\psi_R)$
Q_{G11}	$\bar{\psi}\psi W^A_{\mu\nu} W^A_{\nu\rho} B^\mu B^\rho$	$Q_{\psi^2 X\psi}$	$(\bar{\psi}\psi\psi)_\mu \psi^\mu B_\nu W^{\nu A}$	$Q_{LL}^{(11)}$	$(\bar{\psi}_L\tilde{D}_\mu\psi)(\bar{\psi}_L\psi_L)$	$Q_{LR}^{(11)}$	$(\bar{\psi}_L\tilde{D}_\mu\psi)(\bar{\psi}_R\psi_R)$
Q_{G12}	$\bar{\psi}\psi W^A_{\mu\nu} W^A_{\nu\rho} B^\mu B^\rho$	$Q_{\psi^2 X\psi}$	$(\bar{\psi}\psi\psi)_\mu \psi^\mu B_\nu W^{\nu A}$	$Q_{LL}^{(12)}$	$(\bar{\psi}_L\tilde{D}_\mu\psi)(\bar{\psi}_L\psi_L)$	$Q_{LR}^{(12)}$	$(\bar{\psi}_L\tilde{D}_\mu\psi)(\bar{\psi}_R\psi_R)$
				B-violating			
				$Q_{LL}^{(13)}$	$(\bar{\psi}_L\psi_L)(\bar{\psi}_L\psi_L)$	$Q_{LR}^{(13)}$	$\epsilon^{abc} (\bar{\psi}_L^a\psi_L^b)(\bar{\psi}_L^c\psi_L^c)$
				$Q_{LL}^{(14)}$	$(\bar{\psi}_L\psi_L)(\bar{\psi}_L\psi_L)$	$Q_{LR}^{(14)}$	$\epsilon^{abc} (\bar{\psi}_L^a\psi_L^b)(\bar{\psi}_L^c\psi_L^c)$
				$Q_{LL}^{(15)}$	$(\bar{\psi}_L\psi_L)(\bar{\psi}_L\psi_L)$	$Q_{LR}^{(15)}$	$\epsilon^{abc} (\bar{\psi}_L^a\psi_L^b)(\bar{\psi}_L^c\psi_L^c)$
				$Q_{LL}^{(16)}$	$(\bar{\psi}_L\psi_L)(\bar{\psi}_L\psi_L)$	$Q_{LR}^{(16)}$	$\epsilon^{abc} (\bar{\psi}_L^a\psi_L^b)(\bar{\psi}_L^c\psi_L^c)$
				$Q_{LL}^{(17)}$	$(\bar{\psi}_L\psi_L)(\bar{\psi}_L\psi_L)$	$Q_{LR}^{(17)}$	$\epsilon^{abc} (\bar{\psi}_L^a\psi_L^b)(\bar{\psi}_L^c\psi_L^c)$
				$Q_{LL}^{(18)}$	$(\bar{\psi}_L\psi_L)(\bar{\psi}_L\psi_L)$	$Q_{LR}^{(18)}$	$\epsilon^{abc} (\bar{\psi}_L^a\psi_L^b)(\bar{\psi}_L^c\psi_L^c)$

- ▶ Write down all possible (non-redundant) dimension-6 operators ...
- ▶ **59 operators (76 parameters)** for 1 generation, or **2499 parameters** for 3 generations. [arXiv:1008.4884] Grzadkowski, Iskrzyński, Misiak, Rosiek, [arXiv:1312.2014] Alonso, Jenkins, Manohar, Trott.
- ▶ A **full global fit** with all measurements to all operator coefficients?
 - ▶ We usually only need to deal with a subset of them, e.g. $\sim 20-30$ parameters for **Higgs and electroweak** measurements.
- ▶ Do a global fit and present the results with some fancy bar plots!

You can't really separate Higgs from the EW gauge bosons!

$$\begin{aligned} \mathcal{O}_{H\ell} &= iH^\dagger \overleftrightarrow{D}_\mu H \bar{\ell}_L \gamma^\mu \ell_L, \\ \mathcal{O}'_{H\ell} &= iH^\dagger \sigma^a \overleftrightarrow{D}_\mu H \bar{\ell}_L \sigma^a \gamma^\mu \ell_L, \\ \mathcal{O}_{He} &= iH^\dagger \overleftrightarrow{D}_\mu H \bar{e}_R \gamma^\mu e_R \end{aligned}$$

(or the ones with quarks)

- ▶ modifies gauge couplings of fermions,
- ▶ also generates $hVff$ type contact interaction.



$$\begin{aligned} \mathcal{O}_{HW} &= ig(D^\mu H)^\dagger \sigma^a (D^\nu H) W_{\mu\nu}^a, \\ \mathcal{O}_{HB} &= ig'(D^\mu H)^\dagger (D^\nu H) B_{\mu\nu} \end{aligned}$$

- ▶ generate **aTGCs** $\delta g_{1,Z}$ and $\delta \kappa_\gamma$,
- ▶ also generates **HVV anomalous couplings** such as $hZ_\mu \partial_\nu Z^{\mu\nu}$.



$e^+e^- \rightarrow WW$ with Optimal Observables

▶ TGCs (and additional EFT parameters) are sensitive to the differential distributions!

- ▶ One could do a fit to the binned distributions of all angles.
- ▶ Not the most efficient way of extracting information.
- ▶ Correlations among angles are sometimes ignored.

▶ What are optimal observables?



(See e.g. Z.Phys. C62 (1994) 397-412 Diehl & Nachtmann)

- ▶ In the limit of large statistics (everything is Gaussian) and small parameters (linear contribution dominates), the **best possible reaches** can be derived analytically!


$$\frac{d\sigma}{d\Omega} = S_0 + \sum_i S_{1,i} g_i, \quad c_{ij}^{-1} = \int d\Omega \frac{S_{1,i} S_{1,j}}{S_0} \cdot \mathcal{L},$$

▶ Current work on an improved analysis with machine learning. (Shengdu Chai, JG, Lingfeng Li)

▶ **A more realistic experimental analysis is needed!**




[arXiv:1907.04311] de Blas, Durieux, Grojean, JG, Paul




- ▶ 28-parameter fit projected on Higgs couplings and anomalous triple gauge couplings.
- ▶ $\delta g_H^{ZZ} \approx \delta g_H^{WW}$ from theoretical constraints (gauge invariance & custodial symmetry) and EW measurements.
- ▶ Non-negligible improvement from the 360 GeV run.

SMEFT global fit (Vff couplings) (earlier draft)



- ▶ $U(2)$ symmetry imposed on first two generation quarks.



- ▶ 20-parameter fit (assuming flavor universality in gauge-fermion couplings).
- ▶ See next page for the operator basis.


D6 operators

$\mathcal{O}_H = \frac{1}{2}(\partial_\mu H ^2)^2$	$\mathcal{O}_{GG} = g_s^2 H ^2 G_{\mu\nu}^A G^{A,\mu\nu}$
$\mathcal{O}_{WW} = g^2 H ^2 W_{\mu\nu}^a W^{a,\mu\nu}$	$\mathcal{O}_{y_u} = y_u H ^2 \bar{q}_L H u_R + \text{h.c.} \quad (u \rightarrow t, c)$
$\mathcal{O}_{BB} = g'^2 H ^2 B_{\mu\nu} B^{\mu\nu}$	$\mathcal{O}_{y_d} = y_d H ^2 \bar{q}_L H d_R + \text{h.c.} \quad (d \rightarrow b)$
$\mathcal{O}_{HW} = ig(D^\mu H)^\dagger \sigma^a (D^\nu H) W_{\mu\nu}^a$	$\mathcal{O}_{y_e} = y_e H ^2 \bar{l}_L H e_R + \text{h.c.} \quad (e \rightarrow \tau, \mu)$
$\mathcal{O}_{HB} = ig'(D^\mu H)^\dagger (D^\nu H) B_{\mu\nu}$	$\mathcal{O}_{3W} = \frac{1}{3!} g^{\epsilon abc} W_{\mu\nu}^a W_{\nu\rho}^b W^{c\rho\mu}$
$\mathcal{O}_W = \frac{ig}{2} (H^\dagger \sigma^a \overleftrightarrow{D}_\mu H) D^\nu W_{\mu\nu}^a$	$\mathcal{O}_B = \frac{ig'}{2} (H^\dagger \overleftrightarrow{D}_\mu H) \partial^\nu B_{\mu\nu}$
$\mathcal{O}_{WB} = gg' H^\dagger \sigma^a H W_{\mu\nu}^a B^{\mu\nu}$	$\mathcal{O}_{H\ell} = iH^\dagger \overleftrightarrow{D}_\mu H \bar{\ell}_L \gamma^\mu \ell_L$
$\mathcal{O}_T = \frac{1}{2} (H^\dagger \overleftrightarrow{D}_\mu H)^2$	$\mathcal{O}'_{H\ell} = iH^\dagger \sigma^a \overleftrightarrow{D}_\mu H \bar{\ell}_L \sigma^a \gamma^\mu \ell_L$
$\mathcal{O}_{\ell\ell} = (\bar{\ell}_L \gamma^\mu \ell_L) (\bar{\ell}_L \gamma_\mu \ell_L)$	$\mathcal{O}_{H\bar{e}} = iH^\dagger \overleftrightarrow{D}_\mu H \bar{e}_R \gamma^\mu e_R$
$\mathcal{O}_{Hq} = iH^\dagger \overleftrightarrow{D}_\mu H \bar{q}_L \gamma^\mu q_L$	$\mathcal{O}_{Hu} = iH^\dagger \overleftrightarrow{D}_\mu H \bar{u}_R \gamma^\mu u_R$
$\mathcal{O}'_{Hq} = iH^\dagger \sigma^a \overleftrightarrow{D}_\mu H \bar{q}_L \sigma^a \gamma^\mu q_L$	$\mathcal{O}_{Hd} = iH^\dagger \overleftrightarrow{D}_\mu H \bar{d}_R \gamma^\mu d_R$

- ▶ SILH' basis (eliminate \mathcal{O}_{WW} , \mathcal{O}_{WB} , $\mathcal{O}_{H\ell}$ and $\mathcal{O}'_{H\ell}$)
- ▶ Modified-SILH' basis (eliminate \mathcal{O}_W , \mathcal{O}_B , $\mathcal{O}_{H\ell}$ and $\mathcal{O}'_{H\ell}$) (used here)
- ▶ Warsaw basis (eliminate \mathcal{O}_W , \mathcal{O}_B , \mathcal{O}_{HW} and \mathcal{O}_{HB})


Results from the recent snowmass SMEFT global fit study

[2206.08326] de Blas, Du, Grojean, JG, Miralles, Peskin, Tian, Vos, Vryonidou




Impacts of (lack of) the Z-pole run

[2206.08326] de Blas, Du, Grojean, JG, Miralles, Peskin, Tian, Vos, Vryonidou



- Without good Z-pole measurements, the $eeZh$ contact interaction may have a significant impact on the Higgs coupling determination.
- Current (LEP) Z-pole measurements are not good enough for CEPC Higgs measurements!
- The CEPC Z-pole measurements are!**



What's next?

- ▶ Update measurement inputs (if any)
 - ▶ Any updates to the numbers in CEPC Snowmass report [2205.08553] ?
 - ▶ **This is the most essential part!**

- ▶ More measurements?
 - ▶ Top mass measurement
 - ▶ Diboson measurement ($e^+e^- \rightarrow WW$)
 - ▶ $e^+e^- \rightarrow \gamma\gamma/Z\gamma/ZZ \dots$
 - ▶ ...

- ▶ More interpretations?
 - ▶ Overlap with the new physics white paper?

- ▶ **Timeline/deadline?**

backup slides