

Probing Mirror Twin Higgs with Present and Future Weak Lensing Surveys

Yue-Lin Sming Tsai

(Purple Mountain Observatory)

TOPAC2023

Exploring Mirror Twin Higgs Cosmology with Present and Future Weak Lensing Surveys

Lei Zu^{1, a, b} Chi Zhang^{2, a, b} Hou-Zun Chen,^{a, b} Wei Wang,^{a, b} Yue-Lin Sming Tsai^{3, a, b}
Yuhsin Tsai^{4, c} Wentao Luo,^d Yi-Zhong Fan^{a, b}

^a*Key Laboratory of Dark Matter and Space Astronomy, Purple Mountain Observatory, Chinese Academy of Sciences, Nanjing 210033, China*





^b*School of Astronomy and Space Science, University of Science and Technology of China, Hefei, Anhui 230026, China*

^c*Department of Physics, University of Notre Dame, IN 46556, USA*


^d*Department of Astronomy, School of Physical Sciences, University of Science and Technology of China, Hefei, Anhui 230026, China*

2304.06308

Dark Matter Problems







More and more dSphs were found!



IF GR is correct, it will be difficult to explain the universe without DM assumption.



Fundamental Forces

<i>Strong</i>	 <p>Force which holds nucleus together</p>	<p>Strength</p> <p>1</p>	<p>Range (m)</p> <p>10^{-15} (diameter of a medium sized nucleus)</p>	<p>Particle</p> <p>gluons, π(nucleons)</p>
<i>Electro-magnetic</i>		<p>Strength</p> <p>$\frac{1}{137}$</p>	<p>Range (m)</p> <p>Infinite</p>	<p>Particle</p> <p>photon mass = 0 spin = 1</p>
<i>Weak</i>	 <p>neutrino interaction induces beta decay</p>	<p>Strength</p> <p>10^{-6}</p>	<p>Range (m)</p> <p>10^{-18} (0.1% of the diameter of a proton)</p>	<p>Particle</p> <p>Intermediate vector bosons W^+, W^-, Z_0, mass > 80 GeV spin = 1</p>
<i>Gravity</i>		<p>Strength</p> <p>6×10^{-39}</p>	<p>Range (m)</p> <p>Infinite</p>	<p>Particle</p> <p>graviton ? mass = 0 spin = 2</p>

What is the DM-SM interaction strength?


How is possible that no interaction between $1e-6$ and $1e-39$?

Unless, Gravity is not the fundamental force.




However, all the evidence are all based on gravitational interaction.


Can we see any non-gravitational interaction from gravitational evidence?




Only gravitational interaction?



We shall be able to see non-gravitational interactions from precise cosmological measurements.





N. Craig; adapted from Symmetry Magazine

Mirror Twin Higgs


A solution of the Higgs hierarchy problem.

The hierarchy problem in the SM



- Success of radiative corr. in the SM:

	predicted	observed
top quark	179_{-9}^{+12}	172.7 ± 2.9
Higgs boson	91_{-32}^{+45}	?



- Failure of radiative corr. in Higgs sector:

$$m_h = m_{h_{\text{bare}}} + \delta m_{h,\text{top}} + \dots$$

$$150 = 1354294336587235150 - 1354294336587235000$$

Hierarchy problem:

- 'Conspiracy' to get $m_h \sim M_{\text{EW}} (\ll M_{\text{PL}})$
- Biggest troublemaker is the top quark!

Radiative corrections from top quark

$$\delta m_{h,\text{top}}^2 = -\frac{3}{8\pi^2} \lambda_t^2 \Lambda^2$$


Popular solutions of the Higgs hierarchy problem: SUSY, Mirror Twin Higgs, and so on.

The Hidden Naturalness solution

A concrete example: **Twin Higgs**

Chacko, Goh, Harnik (2005), (up to 10 TeV)



Mirror twin Higgs		
\hat{r}	Flat	$[10^{-3}, 1]$
\hat{v}/v	Flat	$[2, 15]$
$\Delta\hat{N}$	Flat	$[10^{-3}, 1]$



- We only introduce three parameters for a cosmological study.
- DR includes twin neutrinos and photons.


Image credit: Yusin Tsai

The Mirror Twin Higgs



- Matter power spectra are suppressed at a large k region.
- Non-linear effects wash out the DAO features.

Parameter	Prior distribution	Prior range
Cosmology		
$\Omega_b h^2$	Flat	[0.022, 0.023]
$\Omega_{\text{cdm}} h^2$	Flat	[0.112, 0.128]
$100 \cdot \theta_s$	Flat	[1.039, 1.043]
$\ln(A_s \times 10^{10})$	Flat	[2.955, 3.135]
n_s	Flat	[0.941, 0.991]
τ_{reio}	Flat	$[10^{-2}, 0.7]$
Mirror twin Higgs		
\hat{r}	Flat	$[10^{-3}, 1]$
\hat{v}/v	Flat	[2, 15]
$\Delta \hat{N}$	Flat	$[10^{-3}, 1]$
Intrinsic alignment		
A_{IA}	Flat	[-6, 6]
η	Flat	[-6, 6]




$$\theta_s = \frac{r_s(z^*)}{D_A(z^*)}$$

$$r_s(z^*) = \int_{z^*}^{\infty} \frac{dz}{H(z)} c_s(z)$$


$$D_A(z^*) = \int_0^{z^*} \frac{dz}{H(z)}$$

H0 and S8 problem

Constraints on H_0





The LCDM large scale results are smaller H_0 but larger S_8 than the small scale measurements.



L. Perivolaropoulos,
and F. Skara
(2105.05208)


Flat Λ CDM – Growth Tension






Small H_0 but large S_8 ?

If adding a dark radiation component to the Universe, H_0 can be increased while S_8 is still large!



Cosmic Shear
measurement



shape-shap 2 points-correlation: Cosmic Shear



$$\xi_{\pm}^{ij}(\theta) = \langle \epsilon_t^i \epsilon_t^j \rangle \pm \langle \epsilon_x^i \epsilon_x^j \rangle.$$

Image credit:
arXiv:1001.1758



shape-shap 2 points-correlation: Cosmic Shear




$$\xi_{\pm}^{ij}(\theta) = \frac{1}{2\pi} \int_0^{\infty} C_{\text{tot}}^{ij}(\ell) J_{0/4}(\ell \cdot \theta) \ell d\ell$$

$$C_{GG}^{ij}(\ell) = \int_0^{\chi_{z_{\text{max}}}} d\chi \frac{W^i(\chi) W^j(\chi)}{\chi^2} P_{\delta} \left(k = \frac{\ell + 1/2}{\chi}, z(\chi) \right)$$

Telescopes for Cosmic Shear



References: HSC website, Chi Zhang, [Chinese Science Bulletin 66, 1290 (2021)], and 2301.03068



Basic likelihoods:

1. Cosmic shear (DES Y3, shape-shape).
2. CMB (TT, TE, EE, lensing).
3. BAO (BOSS DR12).

Results

When studying how MTH can relax the H_0 and S_8 tensions, we further include two datasets one by one



(iv) The SH0ES likelihood is also a Gaussian distribution with the measurement [74]

$$H_0 = 73.04 \pm 1.04 \text{ km s}^{-1} \text{Mpc}^{-1}.$$

(v) The Planck SZ (2013) likelihood⁴ is described by a Gaussian distribution with the measurement [45]





$$S_8^{\text{SZ}} \equiv \sigma_8 (\Omega_m/0.27)^{0.3} = 0.782 \pm 0.010.$$

The impact of v_{ev} and ΔN



- DES Y3 cannot probe a small scale.
- A large \hat{r} -hat and small ΔN can escape from DES due to mask.





Systematic study



- Non-linear effects for THM are still very similar to Λ CDM.



- The largest discrepancy is from the small scale which is already been masked.

Systematic study





- The shear power spectra computed by HMCode (the IC given by the LCDM) and N-body simulation are similar.
- The main differences are hidden at the small angle region.
- They are hard to statistically distinguished.

MTH with SZ (2013) and SHOES



- DES Y3 strongly disfavour the region of large \hat{r} -hat.
- The future telescopes like CSST can pin down the range of \hat{r} -hat.



MTH can pull the parameter space to a lower S_8 .

Summary

- While the MTH model is presently not a superior solution to the observed H_0 tension compared to the $\Lambda\text{CDM}+\Delta N_{\text{eff}}$ model, we demonstrate that it has the potential to alleviate both the H_0 and S_8 tensions, especially if the S_8 tension.
- The MTH model can relax the tensions while satisfying the DES power spectrum constraint up to $k \sim 10 \text{ h Mpc}^{-1}$.
- We show that the future China Space Station Telescope (CSST) can determine the twin baryon abundance with a 10% level precision.