The 3rd International Workshop on BSM Frontiers: Where to Next?

Contribution ID: 16

## Walking-dilaton hybrid inflation with B - L Higgs embedded in dynamical scalegenesis

Tuesday, 20 August 2024 17:10 (20 minutes)

We propose a hybrid inflationary scenario based on eight-flavor hidden QCD with the hidden colored fermions being in part gauged under  $U(1)_{B-L}$ . This hidden QCD is almost scale-invariant, so-called walking, and predicts the light scalar meson (the walking dilaton) associated with the spontaneous scale breaking, which develops

the Coleman-Weinberg (CW) type potential as the consequence of the nonperturbative

scale anomaly, hence plays the role of an inflaton of the small-field inflation.

The  $U(1)_{B-L}$  Higgs is coupled to the walking dilaton inflaton, which is dynamically induced from the socalled bosonic seesaw mechanism.

We explore the hybrid inflation system involving the walking dilaton inflaton and the  $U(1)_{B-L}$  Higgs as a waterfall field.

We find that observed inflation parameters tightly constrain the  $U(1)_{B-L}$  breaking

scale as well as the walking dynamical scale to be  $\sim 10^9$  GeV and  $\sim 10^{14}$  GeV, respectively, so as to make the waterfall mechanism worked.

The lightest walking pion mass is then predicted to be around 500 GeV. Phenomenological perspectives including embedding of the dynamical electroweak scalegenesis and

possible impacts on the thermal leptogenesis are also addressed.

**Primary authors:** ZHANG, He-Xu (Jilin University); ISHIDA, Hiroyuki (Center for Liberal Arts and Sciences, Toyama Prefectural University, Toyama 939-0398, Japan); LIU, Jie (Jilin University); MATSUZAKI, Shinya (Center for Theoretical Physics and College of Physics, Jilin University, Changchun, 130012, China.)

**Presenter:** LIU, Jie (Jilin University)

Session Classification: Parallel talks (1)