

The near-pericenter self-intersection of the debris stream in TDEs by a misaligned Kerr black hole

Tuesday, 15 October 2024 11:40 (15 minutes)

In TDEs, the stream-stream collision is an important way to dissipate orbital energy in the process of circularizing the debris stream. For non-rotating black holes (BHs), due to the apsidal precession the stream intersects with itself near the its orbital apocenter after a single winding. However, for rotating BHs, the out-of-plane precession arising from the Lense-Thirring effect may prevent the above apocenter intersection, and the stream will instead continue to fall back to pericenter and may self intersect near the pericenter.

We perform local 3-D hydrodynamic simulations of the stream self-intersections near pericenter with Athena++, with the BH's gravity included. The stream components that intersect at the pericenter have a width difference and a vertical offset, thus a large fraction of the stream can avoid the collision. For a BH mass of $10^6 M_{\odot}$ and a orbital penetration factor of 3 and in the case of offset collision where only 14% of the stream collides, we find that 0.5% of the stream kinetic energy is dissipated. After the collision, the stream experiences a lateral expansion, which makes its width to increase with radius more quickly than the case without a pericenter intersection. Our results suggest that for Kerr-BH TDEs which people once thought to have a delayed circularization, the stream intersection at the pericenter is likely to promote collisions in subsequent orbits and accelerate the disc formation.

Primary author: SHEN, Rong-Feng (Sun Yat-Sen University)

Co-author: Ms SONG, Yan (Sun Yat-sen University)

Presenter: SHEN, Rong-Feng (Sun Yat-Sen University)

Session Classification: Session